·Original Article·

# Formaldehyde up-regulates TRPV1 through MAPK and PI3K signaling pathways in a rat model of bone cancer pain

Ying Han<sup>1,\*</sup>, Yan Li<sup>1,\*</sup>, Xing Xiao<sup>1</sup>, Jia Liu<sup>1</sup>, Xiang-Ling Meng<sup>1</sup>, Feng-Yu Liu<sup>1</sup>, Guo-Gang Xing<sup>1</sup>, You Wan<sup>1,2,3</sup>

<sup>1</sup>Neuroscience Research Institute, <sup>2</sup>Department of Neurobiology, School of Basic Medical Sciences, <sup>3</sup>Key Laboratory for Neuroscience, The Ministry of Education / The Ministry of Health, Peking University, Beijing 100191, China

© Shanghai Institutes for Biological Sciences, CAS and Springer-Verlag Berlin Heidelberg 2012

Abstract: Objective Our previous study showed that tumor tissue-derived formaldehyde at low concentrations plays an important role in bone cancer pain through activating transient receptor potential vanilloid subfamily member 1 (TRPV1). The present study further explored whether this tumor tissue-derived endogenous formaldehyde regulates TRPV1 expression in a rat model of bone cancer pain, and if so, what the possible signal pathways are during the development of this type of pain. Methods A rat model of bone cancer pain was established by injecting living MRMT-1 tumor cells into the tibia. The formaldehyde levels were determined by high performance liquid chromatography, and the expression of TRPV1 was examined with Western blot and RT-PCR. In primary cultured dorsal root ganglion (DRG) neurons, the expression of TRPV1 was assessed after treatment with 100 µmol/L formaldehyde with or without pre-addition of PD98059 [an inhibitor for extracellular signal-regulated kinase], SB203580 (a p38 inhibitor), SP600125 [an inhibitor for c-Jun Nterminal kinase], BIM [a protein kinase C (PKC) inhibitor] or LY294002 [a phosphatidylinositol 3-kinase (PI3K) inhibitor]. Results In the rat model of bone cancer pain, formaldehyde concentration increased in blood plasma, bone marrow and the spinal cord. TRPV1 protein expression was also increased in the DRG. In primary cultured DRG neurons, 100 umol/L formaldehyde significantly increased the TRPV1 expression level. Pre-incubation with PD98059, SB203580, SP600125 or LY294002, but not BIM, inhibited the formaldehyde-induced increase of TRPV1 expression. Conclusion Formaldehyde at a very low concentration up-regulates TRPV1 expression through mitogen-activated protein kinase and PI3K, but not PKC, signaling pathways. These results further support our previous finding that TRPV1 in peripheral afferents plays a role in bone cancer pain.

Keywords: formaldehyde; TRPV1; cancer pain; mitogen-activated protein kinase; phosphatidylinositol 3-kinase

## 1 Introduction

Pain is one of the most common symptoms in cancer patients, 75%–90% of advanced or terminal cancer pa-

tients experiencing chronic pain<sup>[1]</sup>. Malignant bone tumors occur in patients with primary bone cancer, but are more common in distant metastases from other primary cancers, notably those of the breast, lung and prostate<sup>[2]</sup>.

Formaldehyde is a consistent metabolic product of demethylation (N-, O-, and S-demethylation)<sup>[3]</sup>. Its concentration is increased in the urine of patients with prostate cancer and bladder cancer and in the expired air from tumor-bearing mice and breast cancer patients<sup>[4,5]</sup> who

<sup>\*</sup>These authors contributed equally to this work.

Corresponding author: You Wan

Tel & Fax: +86-10-82805185

E-mail: ywan@hsc.pku.edu.cn

Article ID: 1673-7067(2012)02-0165-08 Received date: 2011-12-10; Accepted date: 2012-01-27

more frequently suffer from bone cancer pain<sup>[6]</sup>. Formaldehyde is considered a risk factor in the development of chronic bone cancer pain<sup>[7]</sup>, and our recent research demonstrated for the first time that cancer tissue-derived low concentrations of formaldehyde induce cancer pain<sup>[8]</sup>.

Transient receptor potential vanilloid subfamily member 1 (TRPV1), a nociceptive receptor in peripheral nerve fibers<sup>[9]</sup>, plays a pivotal role in the development of cancer pain<sup>[10-16]</sup>. Our previous study found that formaldehyde at concentrations as low as 1 mmol/L induces TRPV1 currents in dorsal root ganglion (DRG) neurons<sup>[8,17]</sup>. It is well-known that TRPV1 expression increases in peripheral nerve fibers and DRG neurons in cancer pain. It is therefore of interest to determine whether formaldehyde modulates TRPV1 expression in peripheral afferents in bone cancer pain, and if so, reveal the intracellular signaling pathway underlying this modulation. Mitogen-activated protein kinases (MAPKs), protein kinase C (PKC) and phosphatidylinositol 3-kinase (PI3K) play critical roles in the cell signaling of TRPV1 regulation. MAPKs have three major family members: extracellular signalregulated kinase (ERK) (ERK1/2 or p44/42 MAPK), p38 and c-Jun N-terminal kinase (JNK), representing three different signaling cascades<sup>[18]</sup>. It was reported that inhibitors of ERK, p38 and JNK alleviate neuropathic pain<sup>[19,20]</sup>, while PKC and PI3K are also involved in the regulation of pain<sup>[20,21]</sup>. These protein kinases, once activated, phosphorylate TRPV1, leading to rapid and dynamic changes in pain sensitivity<sup>[22-26]</sup>. Exposure of intestinal epithelial cells to low-dose formaldehyde results in MAPK activation<sup>[27]</sup>, and formaldehyde induces apoptosis through decreased peroxiredoxin 2 via p38 MAPK in lung epithelial cells<sup>[28]</sup>. So, we set out to determine whether the protein kinase signaling pathway participates in the modulation of TRPV1 expression by formaldehyde in the development of bone cancer pain.

In the present study, a rat model of bone cancer pain and primary cultures of DRG neurons were used to determine whether tumor tissue-derived endogenous formaldehyde regulates TRPV1 expression and if so, the possible underlying signal pathways.

# 2 Materials and methods

2.1 Preparation of MRMT-1 rat mammary gland carcinoma cells MRMT-1 rat mammary gland carcinoma cells were a kind gift from Novartis Oncology Research Institute and were cultured in RPMI-1640 medium (Hyclone, Logan, UT, USA) containing 10% fetal bovine serum (FBS, heat-inactivated) according to our previous description<sup>[8]</sup>. Cells in the culture dish were harvested using 0.25% (*w*/*v*) trypsin (Hyclone), centrifuged and then washed with phosphate-buffered saline (PBS). The final pellet was re-suspended in PBS, and cells were kept on ice until injection into the tibia. In the heat-killed control group, MRMT-1 cells were diluted to the same final concentrations but boiled in water for 20 min before injection. In the PBS control group, only PBS was injected.

2.2 Establishment of the rat model of bone cancer pain Female Sprague-Dawley rats weighing 250-300 g were used. Establishment of the bone cancer pain model has been described previously in detail<sup>[8,29]</sup>. Briefly, after anesthesia, the rat was placed supine. The left hindlimb was shaved and the skin disinfected with 75% (v/v) ethanol. A 1-cm rostro-caudal incision was made in the skin over the top half of the tibia. The tibia was carefully exposed with minimal damage to muscles or blood vessels. Using a 23-gauge needle, the bone was pierced 5 mm below the knee joint distal to the epiphyseal growth plate. The needle was inserted at an angle to enable it to be pushed down the intramedullary canal. Then it was removed and replaced by a long thin blunt needle. A volume of 3  $\mu$ L (3 × 10<sup>4</sup> cells) MRMT-1 live cells, heat-killed MRMT-1 cells or PBS vehicle was injected into the bone cavity. As the plunger in the syringe was slowly depressed, the needle was simultaneously lifted out of the bone to enable the cells to fill the space left in the bone cavity, with no leakage of cells from the bone. Following injection, the site was closed using bone wax. The wound was then closed with suture silk and dusted with penicillin antibiotic powder. The experiments were approved by the Animal Research Committee of Peking University Health Science Center.

**2.3 Primary culture of DRG neurons** Two-week-old rats were decapitated after anesthesia. Lumbar 4 (L4)–L5

DRGs were dissected free. All DRGs on both sides from cervical, thoracic and lumbar levels were collected, placed in a 35-mm culture dish containing 1 mL collagenase (type I, 3 mg/mL; Sigma-Aldrich, St. Louis, MO, USA), and shaken for 45 min in a heated (37°C) chamber. After removing the collagenase and washing with PBS, 1 mL trypsin (0.25%) was added and the dish was shaken for 10 min in the same chamber. Digestion was inhibited by FBS, followed by spinning at 500 rpm for 5 min. Then trypsin was replaced by fresh DMEM containing FBS. The cells were dissociated with a polished Pasteur pipette and plated on poly-*L*-lysine-coated 35-mm plastic dishes or glass sheets for culture.

2.4 Drug administration Formaldehyde was dissolved and diluted in sterile PBS to the desired concentrations (30, 100 and 300 µmol/L). To assess the possible underlying signaling pathways for formaldehyde modulation of TRPV1, 10 µmol/L PD98059 (an ERK inhibitor), 10 µmol/L SB203580 (a p38 inhibitor), 10 µmol/L SP600125 (a JNK inhibitor), 10 µmol/L LY294002 (a PI3K inhibitor) or 1 µmol/L bis-indolylmaleimide I hydrochloride (BIM, a PKC inhibitor) was added to the culture dishes 60 min prior to formaldehyde incubation. All inhibitors were from Sigma-Aldrich Chemicals. Cells without formaldehyde and any inhibitor treatment were used as negative control, and those with formaldehyde but no inhibitor treatment were used as positive control.

**2.5 Measurement of formaldehyde concentration with high performance liquid chromatography (HPLC)** Formaldehyde concentrations in blood plasma, bone marrow, and tissue homogenates were assessed on day 14 following tumor cell inoculation and in the cultural media of cancer cell lines using HPLC (ESA Biosciences, Chelmsford, MA, USA) based on the previous descriptions with modifications<sup>[8,30]</sup>.

**2.6 Western blot detection of TRPV1** L4 and L5 DRGs or primary cultured DRG neurons were dissociated in lysis buffer containing (in mmol/L) 50 Tris-HCl, 40 NaF, 2 EDTA, 1 dithiothreitol and protease inhibitor, pH 7.6. Lysates were separated in SDS-PAGE gel and transferred on to polyvinylidene difluoride membranes.

After blocking with 5% non-fat, dried milk in TBST (20 mmol/L Tris-HCl, pH 7.5, 150 mmol/L NaCl, 0.05% Tween 20), membranes were incubated with rabbit anti-TRPV1 antibody (1:1 000, Abcam, Cambridge, UK) in 5% non-fat, dried milk in TBST overnight at 4°C. After washes with TBST, membranes were incubated with horseradish peroxidase-labeled goat anti-rabbit antibody (Zhongshan Goldenbridge Biotechnology Co., Beijing, China) diluted with 5% non-fat dried milk in TBST and detected with enhanced chemiluminescence reagents (Amersham Biosciences, Arlington Heights, IL, USA). Blots were scanned with Spot Advanced, and the band densities were compared using TotalLab software (version 2.01 Bio-Rad, Hercules, CA, USA). Membrane protein was extracted following the method described previously<sup>[31]</sup>.

2.7 RT-PCR detection of TRPV1 Total RNA was extracted from primary cultured neurons using total RNA kits (Fermentas, Thermo Fisher Scientific, Pittsburgh, PA, USA). First-strand cDNA was synthesized using oligo(dT)<sub>15</sub> primer and M-MuLV reverse transcriptase. PCR products were amplified again using the PCR primers (TRPV1 forward: 5'-GTGCCGGTTTATGTTCGTCT, reverse: 5'-GTT TACCTCGTCCACCCTGA; β-actin forward: 5'-ATCTG GCACCACACCTCC, reverse: 5'-AGCCAGGTCCA GACGCA). PCR products were separated in 2% agarose gel, stained by ethidium bromide, and visualized under UV light. The ratios of optical densities of DNA bands were measured by TotalLab software (version 2.01). All chemicals were from Invitrogen (Grand Island, NY, USA). Cells without formaldehyde and any inhibitor treatment were used as negative control, and those with formaldehyde but no inhibitor treatment as positive control.

**2.8 Data analysis** Data are expressed as mean  $\pm$  SEM. For comparison among groups, ANOVA followed by *post hoc* analysis (Bonferroni test) was used. *P* <0.05 was considered as statistically significant.

#### **3** Results

**3.1 Formaldehyde concentration increased in tibia marrow and tissues from rats with bone cancer pain** Formaldehyde concentrations by HPLC measurement are

Tissues	Control group (µmol/L)	Cancer pain group (µmol/L)
Cerebral cortex (L)	96 ± 17	93 ± 14
Cerebral cortex (R)	$102 \pm 21$	$115 \pm 36$
Hippocampus (L)	$94 \pm 16$	$92 \pm 6$
Hippocampus (R)	$107 \pm 33$	$98 \pm 6$
Thalamus (L)	86 ± 13	91 ± 11
Thalamus (R)	$96 \pm 21$	$96 \pm 11$
Spinal cord (L)	$78 \pm 14$	257 ± 59 **
Spinal cord (R)	$89 \pm 14$	$97 \pm 12$
Bone marrow (L)	$51 \pm 9$	657 ± 104 **
Bone marrow (R)	$54 \pm 15$	96 ± 12 *
Lung	$67 \pm 15$	85 ± 13
Cardiac muscle	$109 \pm 31$	$115 \pm 41$
Pancreas	$75 \pm 14$	$198 \pm 46$ *
Liver	$104 \pm 26$	$174 \pm 102$ *
Spleen	$59 \pm 12$	$100 \pm 23$ *
Kidney	$74 \pm 15$	$101 \pm 16$ *
Blood	$76 \pm 24$	$212 \pm 84$ *

Table 1. Formaldehyde concentrations in blood plasma and tissues

from cancer pain rats compared with control group

Data are expressed as mean  $\pm$  SEM. L, left (ipsilateral) side; R, right side. \*P < 0.05,

\*\*P < 0.01 compared with control. n = 6.

shown in Table 1. In the control group, formaldehyde was detected in all of the collected tissues and blood plasma. Compared with the control group, formaldehyde concentrations significantly increased in the bone cancer pain group on day 14 in plasma, ipsilateral spinal cord, bone marrow, pancreas, liver, spleen and kidney, and especially in bone marrow and ipsilateral spinal cord.

**3.2 Increase of TRPV1 protein expression in DRG in cancer pain rats** Levels of TRPV1 in the DRGs at L4–L5 on days 1, 3, 7, 14, 21 and 28 after inoculation of MRMT-1 tumor cells into the tibia were measured by Western blot. Compared with the PBS and the heat-killed groups, TRPV1 protein expression increased significantly in the bone cancer pain group on day 21, and continued increasing to day 28. In addition, there was no difference in TRPV1 expression between the PBS and heat-killed groups at any of these time points (Fig. 1).

3.3 Formaldehyde treatment increased TRPV1 expression in primary cultured DRG neurons The expression of TRPV1 protein in primary cultured DRG neurons treated with formaldehyde (30, 100, and 300  $\mu$ mol/L) was measured with Western blot at 4, 12, 24, 48 and 72 h of formaldehyde incubation. Compared with the control group, TRPV1 protein increased significantly at 48 h, and



Fig. 1. TRPV1 protein expression increased in lumber (L) 4–L5 dorsal root ganglia in cancer pain rats, as detected by Western blot. A: Western blots of TRPV1 protein on days 1, 3, 7, 14, 21 and 28 following inoculation of MRMT-1 tumor cells into the tibia. P, PBS group; M, MRMT-1 live cells group; H, heat-killed MRMT-1 cells group. B: Statistical analysis of the relative densities of TRPV1 bands. Compared with the heat-killed or PBS group, TRPV1 protein increased significantly on day 21, and continued increasing to day 28 after inoculation of MRMT-1 cancer cells. \*P <0.05, \*\*P < 0.01 compared with control. n = 6 rats for each time point in each group.</li>



Fig. 2. Increase of transient receptor potential vanilloid subfamily member 1 (TRPV1) expression in primary cultured dorsal root ganglion neurons after formaldehyde incubation. A: Example of TRPV1 protein bands detected by Western blot. B: Statistical analysis of the relative amount of TRPV1 protein. C: Example of TRPV1 mRNA bands detected by RT-PCR. D: Statistical analysis of the relative amount of TRPV1 mRNA. Compared with the PBS control group, TRPV1 expression increased at 48 h of 100 µmol/L formaldehyde treatment, and further increased at 72 h. \**P* <0.05, \*\**P* <0.01 compared with control (*n* = 6).

further increased at 72 h of incubation in the 100  $\mu$ mol/L formaldehyde group (Fig. 2A, B). Besides, there was no difference in TRPV1 protein expression between the groups treated with 30 and 300  $\mu$ mol/L formaldehyde and the control group (data not shown).

In addition, TRPV1 mRNA levels detected by RT-PCR showed changes consistent with those of protein expression. In the 100  $\mu$ mol/L formaldehyde group, TRPV1 mRNA increased at 48 h, and further increased at 72 h (Fig. 2C, D). Similarly, the expression of TRPV1 mRNA in the groups treated with 30  $\mu$ mol/L and 300  $\mu$ mol/L formaldehyde did not differ from that in the control group (data not shown).

**3.4 Inhibitory effects of MAPK and PI3K inhibitors on formaldehyde-induced TRPV1 up-regulation in primary cultured DRG neurons** After 72-h incubation of cultured DRG neurons with 100 μmol/L formaldehyde alone, TRPV1 protein expression increased significantly (Fig. 3). Compared with the formaldehyde-only group, the TRPV1

expression decreased after pre-incubation with the signal transduction pathway inhibitors PD98059, SB203580, SP600125 or LY294002. However, TRPV1 protein expression did not change after addition of BIM (Fig. 3A, B).

Furthermore, changes of TRPV1 mRNA expression in DRG neurons were consistent with those of TRPV1 protein expression. TRPV1 mRNA increased significantly 72 h after 100  $\mu$ mol/L formaldehyde treatment. Compared with the formaldehyde-only group, TRPV1 mRNA expression decreased in the formaldehyde groups with pre-addition of the inhibitors PD98059, SB203580, SP600125 or LY294002, but did not change after addition of BIM (Fig. 3C, D).

#### 4 Discussion

Formaldehyde is a consistent metabolic product of demethylation. Previous studies demonstrated that the formaldehyde level increases in some tissues and the blood of patients with bone cancer pain, but formaldehyde is



Fig. 3. Effects of signal transduction pathway inhibitors on formaldehyde (FA)-induced up-regulation of transient receptor potential vanilloid subfamily member 1 (TRPV1) expression in primary cultured dorsal root ganglion (DRG) neurons. A: Example of TRPV1 protein bands detected by Western blot. B: Statistical analysis of the relative band densities of TRPV1 protein. C: Example of TRPV1 mRNA bands detected by RT-PCR. D: Statistical analysis of the relative band densities of TRPV1 mRNA. TRPV1 expression increased significantly in DRG neurons 72 h after incubation with 100 μmol/L formaldehyde. Compared with formaldehyde alone, PD98059 (inhibitor for extracellular signal-regulated kinase), SB203580 (p38 inhibitor), SP600125 (inhibitor for c-Jun *N*-terminal kinase), and LY294002 (inhibitor for phosphatidylinositol 3-kinase) inhibited the formaldehyde-induced up-regulation of TRPV1 expression. BIM (inhibitor for protein kinase C) had no significant effect. \**P* <0.01 compared with control, \**P* <0.05 compared with formaldehyde group (*n* = 6).

usually considered as a risk factor (a cause) in the development of cancer<sup>[4-7]</sup>. In the present study, formaldehyde concentrations in blood and tissues increased significantly in rats with bone cancer pain. This result is consistent with our previous finding that formaldehyde is increased in the cancer tissues of patients<sup>[8]</sup>. More importantly, this result in a rat model of cancer pain provides further support for the concept that the elevated formaldehyde is not only a "cause", but also, more interestingly, a "consequence" of the pathogenesis of cancer or bone cancer pain.

TRPV1 is expressed predominantly in peripheral nociceptors and participates in pain sensitization<sup>[32]</sup>. Our previous study found that cancer tissue-derived endogenous formaldehyde at low pathological concentrations in an acidic environment induces pain responses through TRPV1 activation in rats *in vivo*<sup>[8]</sup>. In the present study, our

results showed that TRPV1 protein expression increased in rats with bone cancer pain, and formaldehyde incubation increased TRPV1 expression in cultured DRG neurons. Here, formaldehyde incubation with cultured DRG neurons mimics the contact of cancer tissue-derived formaldehyde with peripheral afferent nerve fibers innervating the tibia in the rat model of bone cancer pain. These results indicate that formaldehyde induces bone cancer pain not only by activating TRPV1 receptors (as in our previous report<sup>[8]</sup>), but also by up-regulating TRPV1 protein expression in peripheral nociceptors.

It is an interesting phenomenon that endogenous formaldehyde at very low concentrations up-regulates TRPV1 expression. To further investigate the mechanisms underlying this, we examined possible signaling transduction pathways involved in the process. The results showed that application of MAPK and PI3K inhibitors suppressed the formaldehyde-induced increase of TRPV1 expression in cultured DRG neurons, while a PKC inhibitor did not. It is well known that MAPKs, PI3K and PKC play critical roles in cell signaling. MAPKs have three major family members, ERK, p38, and JNK, that represent three different signaling cascades<sup>[33]</sup>. TRPV1 expression can result from ERK and p38 signaling<sup>[34,35]</sup>. PI3K is another key mediator of central pain sensitization and of inflammatory heat hyperalgesia through TRPV1 sensitization<sup>[24,36]</sup>. So, the present study gives preliminary evidence that formaldehyde up-regulates TRPV1 expression through MAPK and PI3K signal pathways, but not the PKC signal pathway. However, more evidence is needed and related interesting questions, for example, interaction or cross-talk among these signaling pathways, are worthy of further investigation.

Previous studies showed that MAPKs, PI3K and their downstream kinases in primary sensory neurons are involved in pain<sup>[20,21]</sup>. MAPK and PI3K pathways are usually targets of activating factors in the tumor microenvironment, and increased formaldehyde activates MAPK and PI3K/AKT signal pathways<sup>[37]</sup>. PI3K in primary DRG sensory neurons is activated by inflammatory agents, and PI3K inhibitors inhibit chemical- and nerve injury-induced pain hypersensitivity, suggesting that PI3K activation contributes to pain<sup>[24,38]</sup>.

We note possible differences between the responses of DRG neurons to formaldehyde in two-week-old rat culture and those in adult animals. We chose two-week-old culture because long-term culture of the adult DRG is technically very difficult.

In summary, on the basis of our previous report that tumor tissue-derived endogenous formaldehyde at very low concentrations induces cancer pain through activation of TRPV1, the present study provides further evidence that formaldehyde at a low pathological concentration up-regulates TRPV1 expression through MAPK and PI3K signaling pathways, but not the PKC pathway, in the pathogenesis of bone cancer pain.

Acknowledgements: This work was supported by

grants from the National Natural Science Foundation of China (81070893, 81171042 and 31171063), Beijing Municipal Commission of Education "Grants for Outstanding Ph.D. Program Tutors", the "111" Project of the Ministry of Education of China, and The Ministry of Education (BMU20100014) and the China Postdoctoral Science Foundation (20090450266).

## **References:**

- Portenoy RK, Payne D, Jacobsen P. Breakthrough pain: characteristics and impact in patients with cancer pain. Pain 1999, 81: 129–134.
- [2] Ghilardi JR, Röhrich H, Lindsay TH, Sevcik MA, Schwei MJ, Kubota K, et al. Selective blockade of the capsaicin receptor TRPV1 attenuates bone cancer pain. J Neurosci 2005, 25: 3126– 3131.
- [3] Kalasz H. Biological role of formaldehyde, and cycles related to methylation, demethylation, and formaldehyde production. Mini Rev Med Chem 2003, 3: 175–192.
- [4] Ebeler SE, Clifford AJ, Shibamoto T. Quantitative analysis by gas chromatography of volatile carbonyl compounds in expired air from mice and human. J Chromatogr B Biomed Sci Appl 1997, 702: 211–215.
- [5] Spanel P, Smith D, Holland TA, Al Singary W, Elder JB. Analysis of formaldehyde in the headspace of urine from bladder and prostate cancer patients using selected ion flow tube mass spectrometry. Rapid Commun Mass Spectrom 1999, 13: 1354–1359.
- [6] Sabino MA, Mantyh PW. Pathophysiology of bone cancer pain. J Support Oncol 2005, 3: 15–24.
- [7] Thorndike J, Beck WS. Production of formaldehyde from N5methyltetrahydrofolate by normal and leukemic leukocytes. Cancer Res 1977, 37: 1125–1132.
- [8] Tong Z, Luo W, Wang Y, Yang F, Han Y, Li H, et al. Tumor tissuederived formaldehyde and acidic microenvironment synergistically induce bone cancer pain. PLoS One 2010, 5: e10234.
- [9] Pei L, Lin CY, Dai JP, Yin GF. Facial pain induces the alteration of transient receptor potential vanilloid receptor 1 expression in rat trigeminal ganglion. Neurosci Bull 2007, 23: 92–100.
- [10] Luo H, Cheng J, Han JS, Wan Y. Change of vanilloid receptor 1 expression in dorsal root ganglion and spinal dorsal horn during inflammatory nociception induced by complete Freund's adjuvant in rats. Neuroreport 2004, 15: 655–658.
- [11] Christoph T, Grunweller A, Mika J, Schäfer MK, Wade EJ, Weihe E, et al. Silencing of vanilloid receptor TRPV1 by RNAi reduces neuropathic and visceral pain *in vivo*. Biochem Biophys Res Commun

2006, 350: 238-243.

- [12] Niiyama Y, Kawamata T, Yamamoto J, Omote K, Namiki A. Bone cancer increases transient receptor potential vanilloid subfamily 1 expression within distinct subpopulations of dorsal root ganglion neurons. Neuroscience 2007, 148: 560–572.
- [13] Yu L, Yang F, Luo H, Liu FY, Han JS, Xing GG, et al. The role of TRPV1 in different subtypes of dorsal root ganglion neurons in rat chronic inflammatory nociception induced by complete Freund's adjuvant. Mol Pain 2008, 4: 61.
- [14] Niiyama Y, Kawamata T, Yamamoto J, Furuse S, Namiki A. SB366791, a TRPV1 antagonist, potentiates analgesic effects of systemic morphine in a murine model of bone cancer pain. Br J Anaesth 2009, 102: 251–258.
- [15] Du Y, Xiao Y, Lu ZM, Ding J, Xie F, Fu H, *et al.* Melittin activates TRPV1 receptors in primary nociceptive sensory neurons via the phospholipase A2 cascade pathways. Biochem Biophys Res Commun 2011, 408: 32–37.
- [16] Xia R, Samad TA, Btesh J, Jiang LH, Kays I, Stjernborg L, et al. TRPV1 signaling: mechanistic understanding and therapeutic potential. Curr Top Med Chem 2011, 11: 2180–2191.
- [17] Tian LJ, Du Y, Xiao Y, Lv ZM, Yu YQ, Cui XY, et al. Mediating roles of the vanilloid receptor TRPV1 in activation of rat primary afferent nociceptive neurons by formaldehyde. Acta Physiol Sin 2009, 61: 404–416.
- [18] Svensson C, Part K, Kunnis-Beres K, Kaldmäe M, Fernaeus SZ, Land T. Pro-survival effects of JNK and p38 MAPK pathways in LPS-induced activation of BV-2 cells. Biochem Biophys Res Commun 2011, 406: 488–492.
- [19] Doya H, Ohtori S, Fujitani M, Saito T, Hata K, Ino H, et al. c-Jun N-terminal kinase activation in dorsal root ganglion contributes to pain hypersensitivity. Biochem Biophys Res Commun 2005, 335: 132–138.
- [20] Ji RR, Kawasaki Y, Zhuang ZY, Wen YR, Zhang YQ. Protein kinases as potential targets for the treatment of pathological pain. Handb Exp Pharmacol 2007: 359–389.
- [21] Ji RR, Strichartz G. Cell signaling and the genesis of neuropathic pain. Sci STKE 2004, 2004: E14.
- [22] Ji RR, Samad TA, Jin SX, Schmoll R, Woolf CJ. p38 MAPK activation by NGF in primary sensory neurons after inflammation increases TRPV1 levels and maintains heat hyperalgesia. Neuron 2002, 36: 57–68.
- [23] Morenilla-Palao C, Planells-Cases R, Garcia-Sanz N, Ferrer-Montiel A. Regulated exocytosis contributes to protein kinase C potentiation of vanilloid receptor activity. J Biol Chem 2004, 279: 25665–25672.
- [24] Zhuang ZY, Xu H, Clapham DE, Ji RR. Phosphatidylinositol 3-kinase activates ERK in primary sensory neurons and mediates inflammatory heat hyperalgesia through TRPV1 sensitization. J

Neurosci 2004, 24: 8300-8309.

- [25] Amadesi S, Cottrell GS, Divino L, Chapman K, Grady EF, Bautista F, et al. Protease-activated receptor 2 sensitizes TRPV1 by protein kinase C epsilon- and A-dependent mechanisms in rats and mice. J Physiol 2006, 575(Pt 2): 555–571.
- [26] Zhang H, Cang CL, Kawasaki Y, Liang LL, Zhang YQ, Ji RR, et al. Neurokinin-1 receptor enhances TRPV1 activity in primary sensory neurons via PKC epsilon: a novel pathway for heat hyperalgesia. J Neurosci 2007, 27: 12067–12077.
- [27] Feick P, Haas SR, Singer MV, Böcker U. Low-dose exposure of intestinal epithelial cells to formaldehyde results in MAP kinase activation and molecular alteration of the focal adhesion protein paxillin. Toxicology 2006, 219: 60–72.
- [28] Lim SK, Kim JC, Moon CJ, Kim GY, Han HJ, Park SH. Formaldehyde induces apoptosis through decreased Prx 2 via p38 MAPK in lung epithelial cells. Toxicology 2010, 271: 100–106.
- [29] Medhurst SJ, Walker K, Bowes M, Kidd BL, Glatt M, Muller M, et al. A rat model of bone cancer pain. Pain 2002, 96: 129–140.
- [30] Fujita M, Ueda T, Handa T. Generation of formaldehyde by pharmaceutical excipients and its absorption by meglumine. Chem Pharm Bull (Tokyo) 2009, 57: 1096–1099.
- [31] O'Neil KA, Miller FR, Barder TJ, Lubman DM. Profiling the progression of cancer: separation of microsomal proteins in MCF10 breast epithelial cell lines using nonporous chromatophoresis. Proteomics 2003, 3: 1256–1269.
- [32] Wang Y. The functional regulation of TRPV1 and its role in pain sensitization. Neurochem Res 2008, 33: 2008–2012.
- [33] Cheng JK, Ji RR. Intracellular signaling in primary sensory neurons and persistent pain. Neurochem Res 2008, 33: 1970–1978.
- [34] Bron R, Klesse LJ, Shah K, Parada LF, Winter J. Activation of Ras is necessary and sufficient for upregulation of vanilloid receptor type 1 in sensory neurons by neurotrophic factors. Mol Cell Neurosci 2003, 22: 118–132.
- [35] Chen Y, Geis C, Sommer C. Activation of TRPV1 contributes to morphine tolerance: involvement of the mitogen-activated protein kinase signaling pathway. J Neurosci 2008, 28: 5836–5845.
- [36] Pezet S, Marchand F, D'Mello R, Grist J, Clark AK, Malcangio M, *et al.* Phosphatidylinositol 3-kinase is a key mediator of central sensitization in painful inflammatory conditions. J Neurosci 2008, 28: 4261–4270.
- [37] Bonnas C, Specht K, Spleiss O, Froehner S, Dietmann G, Krüger JM, et al. Effects of cold ischemia and inflammatory tumor microenvironment on detection of PI3K/AKT and MAPK pathway activation patterns in clinical cancer samples. Int J Cancer 2011 [Epub ahead of print]
- [38] Shi TJ, Huang P, Mulder J, Ceccatelli S, Hokfelt T. Expression of p-Akt in sensory neurons and spinal cord after peripheral nerve injury. Neurosignals 2009, 17: 203–212.