

AMPA receptor trafficking in inflammation-induced dorsal horn central sensitization

Yuan-Xiang Tao

Department of Anesthesiology and Critical Care Medicine, Johns Hopkins University School of Medicine, Baltimore, Maryland 21205, USA

© Shanghai Institutes for Biological Sciences, CAS and Springer-Verlag Berlin Heidelberg 2012

Abstract: Activity-dependent postsynaptic receptor trafficking is critical for long-term synaptic plasticity in the brain, but it is unclear whether this mechanism actually mediates the spinal cord dorsal horn central sensitization (a specific form of synaptic plasticity) that is associated with persistent pain. Recent studies have shown that peripheral inflammation drives changes in α -amino-3-hydroxy-5-methyl-4-isoxazolepropionic acid receptor (AMPA) subunit trafficking in the dorsal horn and that such changes contribute to the hypersensitivity that underlies persistent pain. Here, we review current evidence to illustrate how spinal cord AMPARs participate in the dorsal horn central sensitization associated with persistent pain. Understanding these mechanisms may allow the development of novel therapeutic strategies for treating persistent pain.

Keywords: dorsal horn; GluA1; GluA2; inflammation; persistent pain; receptor trafficking

1 Introduction

Persistent or chronic pain arising from inflammation and tissue or nerve injury is one of the most common syndromes seen clinically. Despite intensive research into the neurobiologic mechanisms of persistent pain during the past decades, it is still poorly managed by current drugs such as opioids and nonsteroidal anti-inflammatory drugs^[1]. Persistent pain is characterized by spontaneous or intermittent burning pain, an exaggerated response to painful stimuli (hyperalgesia), and pain in response to normally innocuous stimuli (allodynia). It is believed that a specific form of synaptic plasticity known as central sensitization underlies the mechanisms by which persistent pain develops and is maintained. Understanding such mechanisms will

allow the development of novel therapeutic strategies to treat persistent pain.

The α -amino-3-hydroxy-5-methyl-4-isoxazolepropionic acid (AMPA) receptor (AMPA) is a glutamate ionotropic receptor that is integral to fast excitatory synaptic transmission and plasticity at synapses in the central nervous system^[2,3]. In addition to mediating the acute spinal processing of nociceptive and non-nociceptive inputs, AMPARs in the dorsal horn may also participate in central sensitization under persistent pain conditions^[4]. Multiple intrathecal treatment with the non-selective AMPAR antagonist CNQX beginning 15 min prior to nerve ligation significantly (but not completely) reduces the development of nerve injury-induced thermal hyperalgesia, and this effect persists for up to 10 days^[5]. Interestingly, single intrathecal post-treatment with selective or nonselective AMPAR antagonists partially and temporarily attenuates thermal hyperalgesia and mechanical allodynia after peripheral nerve injury^[5,6]. Inhibition of spinal Ca^{2+} -permeable AMPAR blocks thermal

Corresponding author: Yuan-Xiang Tao
Tel: +1-443-2875490; Fax: +1-410-5025554
E-mail: ytao1@jhmi.edu; ytau@jhmi.edu
Article ID: 1673-7067(2012)02-0111-10
Received date: 2012-01-06; Accepted date: 2012-01-27

injury-induced mechanical allodynia and carrageenan-induced thermal hyperalgesia and mechanical allodynia^[7,8]. Furthermore, intrathecal administration of selective AMPAR antagonists attenuates complete Freund's adjuvant (CFA)-induced mechanical and thermal pain hypersensitivity during both the development and maintenance of persistent pain at doses that do not affect locomotor function or responsiveness to acute pain^[9]. Electrophysiologic studies revealed that AMPAR blockade reduces the receptive field areas of superficial dorsal horn neurons^[10,11]. Using mice that lack GluA1, GluA2, or GluA3, Hartmann *et al.*^[12] demonstrated that AMPARs are critical determinants of spinal cord nociceptive plasticity and inflammatory pain. The evidence suggests that dorsal horn AMPARs may be critical for the central sensitization associated with persistent pain. However, how spinal AMPARs participate in central sensitization under persistent pain conditions is still incompletely understood.

Recent studies have indicated that peripheral inflammatory insults drive changes in AMPAR subunit trafficking via N-methyl-D-aspartic acid (NMDA) receptor-triggered activation of protein kinases in the dorsal horn and that these changes contribute to central sensitization in the development and maintenance of persistent inflammatory pain^[4]. Here I review the current evidence regarding changes in AMPAR subunit trafficking in dorsal horn neurons after peripheral inflammation and tissue or nerve injury and discuss potential mechanisms by which such changes participate in the development and maintenance of persistent pain.

2 AMPAR channel structure and expression in spinal cord

AMPARs are composed of four subunits, GluA1 to GluA4 (or GluR1 to GluR4). Each subunit comprises an N-terminal extracellular amino domain, a ligand-binding domain, a receptor-channel domain, and an intracellular C-terminal domain^[3]. It should be emphasized that the receptor-channel domain consists of three transmembrane segments (M1, M3, and M4) and one re-entrant loop within the membrane (M2). Because the M2 loop forms part of

the channel pore, it controls the flow of ions (e.g. Ca^{2+}) through the AMPAR channel^[13,14]. In addition, the C-terminal intracellular domain presents multiple protein phosphorylation sites for various known protein kinases such as protein kinase A (PKA), protein kinase C (PKC), and Ca^{2+} /calmodulin-dependent protein kinase II (CaMKII), and several binding motifs for various membrane-related and intracellular proteins, e.g. the scaffolding proteins^[13,14]. Functional AMPARs are homomeric or heteromeric tetramers of GluR subunits. In the adult mammalian central nervous system, GluA1 and GluA2 subunits are ubiquitously expressed and present in most AMPARs. Whereas GluA1, GluA3, and GluA4 convey high Ca^{2+} permeability to AMPAR channels, GluA2 imparts low Ca^{2+} permeability because it carries an arginine residue in its pore-forming M2 segment (inserted by Q/R site RNA editing during development)^[3].

Morphological studies indicate that all four GluR subunits are found within the dorsal horn of the spinal cord^[15–17]. GluA1 is highly expressed in the superficial dorsal horn (laminae I–II) but exhibits low expression in the deeper dorsal horn^[4]. GluA2 expression has been observed throughout the dorsal horn, but it is distributed predominantly in inner lamina II^[4]. GluA3 is weakly expressed in the deep laminae. GluA4 was recently detected in large NK1 receptor-expressing projection neurons in lamina I^[18]. Under electron microscopy, most GluA1 and GluA2 immunoreactive products are restricted to the postsynaptic density, with little reactivity in axons and terminals in the superficial dorsal horn^[4]. Thus, GluA1 and GluA2 are the two key subunits that comprise the AMPAR channel in the dorsal horn.

3 Trafficking and functional changes of dorsal horn AMPARs after peripheral noxious stimuli

Unlike the NMDA receptor, the AMPAR is not stable at synapses; its expression is dependent on a balance between two dynamic processes. In one, AMPAR subunits rapidly and constitutively cycle between intracellular stores and the cellular membrane surface. In the second, AMPAR subunits exchange between the extrasynaptic and

synaptic membranes by lateral diffusion. Any factors that disturb these processes can change the synaptic expression of AMPAR subunits and further affect AMPAR synaptic strength.

Recent studies have demonstrated that peripheral noxious insults upset the flux of AMPAR subunits and alter their synaptic and extrasynaptic trafficking in dorsal horn neurons. One study showed that painful visceral stimulation produced by capsaicin injection into the colon rapidly and significantly increases the level of membrane GluA1 protein and correspondingly reduces the amount of cytosolic GluA1, without changing total GluA1 or GluA2 protein levels in the dorsal horn^[19] (Table 1). Another study demonstrated that capsaicin injection into rat hind-paw increases the density of GluA1-containing AMPARs and the ratio of GluA1 to GluA2/3 in postsynaptic membrane in the superficial dorsal horn^[20]. In addition, pain hypersensitivity evoked by intraplantar carrageenan and pain behaviors induced by intraplantar formalin are associated with the movement of GluA1, but not GluA2, into neuronal membrane in the dorsal horn^[21,22] (Table 1). Interestingly, intraplantar injection of CFA into a hind-paw produces changes in the subcellular distribution of both GluA1 and GluA2 proteins in dorsal horn neurons, although it does not alter the total expression of GluA1 and GluA2 proteins^[9,23] (Table 1). The level of cytosolic GluA2 is markedly increased, and the level of membrane-bound GluA2 is significantly decreased in the dorsal horn 1 and 3 days after CFA injection^[9,23].

Table 1. Trafficking changes of GluA1 and GluA2 in dorsal horn after inflammation

| Model | GluA1 | | | GluA2 | | |
|---------------------------------|---------|----------|-------|---------|----------|-------|
| | Cytosol | Membrane | Total | Cytosol | Membrane | Total |
| ICo capsaicin ^[19] | ↓ | ↑ | — | — | — | — |
| IPI formalin ^[22] | ↓↑ | ↑ | N/A | N/A | N/A | N/A |
| IPI carrageenan ^[21] | ↓ | ↑ | N/A | — | — | N/A |
| IPI CFA ^[9,23] | ↓ | ↑ | — | ↑ | ↓ | — |

CFA, complete Freund's adjuvant; ICo, intracolonic; IPI, intraplantar; N/A, not available; “↓”, decrease; “↑”, increase; “↓↑”, decrease early and increase late; “—”, no change.

Conversely, the amount of GluA1 is significantly reduced in cytosol and increased in the neuronal membrane 1 day after CFA injection^[9]. Furthermore, researchers found decreased synaptic GluA2 and increased extrasynaptic GluA1 in the superficial dorsal horn 1 day after CFA injection^[23-25]. In contrast, peripheral incision and nerve injury failed to alter the expression and localization of GluA1 or GluA2 in cytosolic and plasma membrane fractions from the dorsal horn^[26,27]. This evidence indicates that different peripheral nociceptive insults produce distinct changes in dorsal horn AMPAR subunit trafficking.

Peripheral noxious insults also alter the properties of synaptic and extrasynaptic AMPARs in dorsal horn neurons. One key property that is altered is Ca^{2+} permeability. Selectively blocking Ca^{2+} -permeable AMPARs inhibits AMPAR-mediated evoked excitatory postsynaptic currents (eEPSCs) to a greater degree in lamina II neurons from CFA-treated animals than in those from saline-treated animals^[23], indicating that CFA injection increases synaptic Ca^{2+} -permeable AMPARs in the superficial dorsal horn. Compared to saline, CFA also enhances the inward rectification of eEPSCs and significantly reduces the rectification index in lamina I projection neurons^[28] and lamina II interneurons^[23,24] 1 day after CFA injection. In addition, extrasynaptic AMPAR currents and Ca^{2+} permeability increase dramatically in the lamina II neurons characterized by intrinsic tonic firing properties, but not in those that exhibit strong adaptation, 1 day after CFA injection^[25]. This increase is accompanied by an inward rectification of AMPA-induced currents and enhanced sensitivity to a highly selective Ca^{2+} -permeable AMPAR blocker^[25]. The Ca^{2+} -permeability of AMPARs can also be visualized by kainate-induced cobalt uptake. One day after injecting one hind-paw of rats with CFA, we found that the number of cobalt-positive neurons was significantly increased in dorsal horn neurons on the side ipsilateral to the injection^[25]. Taken together, these findings suggest that peripheral inflammation leads to a switch from Ca^{2+} -impermeable AMPARs to Ca^{2+} -permeable AMPARs at both the synaptic and extrasynaptic sites of dorsal horn neurons. This switch is very likely attributable to a change in AMPAR subunit

composition (that is, GluA1 membrane insertion and GluA2 internalization).

4 Molecular mechanisms that underlie inflammation-induced changes in dorsal horn AMPAR trafficking

As noted above, the C-terminal intracellular domain of AMPAR subunits contains multiple protein phosphorylation sites for various protein kinases and several binding motifs for scaffolding proteins. New research has revealed that these phosphorylation sites and binding motifs impact AMPAR subunit trafficking and function. For example, phosphorylation of GluA1 at the site of serine 845 (Ser⁸⁴⁵) by PKA elevates the channel open probability and peak amplitude of the AMPAR, increases GluA1 expression in the plasma membrane, and reduces AMPAR internalization^[29–31]. Phosphorylation of GluA1 at Ser⁸³¹ by PKC/CaMKII increases the signal channel conductance of synaptic AMPARs^[32]. CaMKII activation also drives GluA1-containing AMPARs to synapses by a mechanism that requires the binding of GluA1 to its scaffolding protein^[33]. In addition, phosphorylation of GluA1 Ser⁸¹⁸ by PKC promotes GluA1 membrane insertion and facilitates an interaction with protein 4.1N, a scaffolding protein involved in the synaptic delivery and stabilization of GluA1^[34,35]. Further study showed that protein 4.1N is required for activity-dependent GluA1 insertion at extrasynaptic sites by PKC-dependent phosphorylation of GluA1 Ser⁸¹⁶ and Ser⁸¹⁸. Phosphorylation at these sites enhances 4.1N binding to GluA1 and facilitates GluA1-containing AMPAR insertion into the extrasynaptic plasma membrane^[36]. Thus, it is very likely that protein 4.1N and several serine phosphorylation sites of the GluA1 C-terminus participate in the regulation of synaptic and extrasynaptic GluA1 trafficking.

Peripheral inflammation alters the phosphorylation status of GluA1 serine residues. Intraplantar injection of capsaicin or CFA increases the phosphorylation of GluA1 Ser⁸³¹ and Ser⁸⁴⁵ in the dorsal horn^[37–39]. Our recent data demonstrated that dorsal horn PKC α knockdown blocks CFA-induced increases in the number of Ca²⁺-permeable AMPARs and the proportion of Ca²⁺-permeable subunits

at the extrasynaptic site. From these data, it appears that the inflammation-induced increase in extrasynaptic Ca²⁺-permeable AMPARs is due to PKC-dependent phosphorylation of GluA1 at the serine residues and subsequent insertion of GluA1-containing AMPARs into the extrasynaptic plasma membrane of dorsal horn neurons. However, this conclusion must be further confirmed by investigating whether targeted mutation of GluA1 phosphorylation sites (e.g., Ser⁸³¹, Ser⁸⁴⁵, Ser⁸¹⁶ and Ser⁸¹⁸) affects inflammation-induced increases in GluA1 membrane insertion and the number of Ca²⁺-permeable AMPARs at extrasynaptic sites of dorsal horn neurons.

Recent evidence has revealed that spinal cord glial activation is involved in the increase of GluA1 phosphorylation and changes in GluA1 trafficking in the dorsal horn after inflammation. Choi *et al.*^[21] showed that intraplantar carrageenan promoted GluA1 membrane insertion and phosphorylation of GluA1 Ser⁸⁴⁵ in the dorsal horn but that these increases were blocked by a tumor necrosis factor (TNF) antagonist. Thus, the increased GluA1 phosphorylation and membrane insertion may be dependent on TNF, a proinflammatory cytokine. Additional investigation is needed to determine how TNF released from spinal glia participates in the regulation of neuronal GluA1 phosphorylation and trafficking in the dorsal horn after inflammation.

Like GluA1, GluA2 contains a binding motif at its C-terminus for two scaffolding proteins: AMPAR-binding protein (ABP)/glutamate receptor-interacting protein (GRIP) and protein interacting with C kinase 1 (PICK1). ABP/GRIP functions as an anchor to stabilize GluA2 at the synaptic membrane^[40–42]. PICK1 acts as a transporter to recruit PKC α to ABP/GRIP-GluA2 complexes; the kinase then phosphorylates GluA2 at Ser⁸⁸⁰^[43–45]. Studies from *in vitro* cell lines and neuronal culture have shown that GluA2 phosphorylation at Ser⁸⁸⁰ by PKC α reduces the binding of GluA2 to ABP/GRIP, but not to PICK1, and promotes GluA2 internalization^[43,44,46,47].

The data from our laboratory indicate that PKC phosphorylation at GluA2 Ser⁸⁸⁰ is triggered by NMDA receptor activation and contributes to dorsal horn GluA2 inter-

nalization in inflammatory pain. Under normal conditions, dorsal horn NMDA receptors are inactive. GluA2 interacts with both ABP/GRIP and PICK1, but PICK1 also binds to inactive PKC α in dorsal horn neurons^[23]. In cultured dorsal horn neurons, activation of NMDA receptors and PKC α markedly increases the level of GluA2 phosphorylation at Ser⁸⁸⁰ and decreases the surface expression of GluA2^[23]. Activation of AMPARs and group 1 metabotropic glutamate receptors does not have this effect. Consistently, blockade of dorsal horn NMDA receptors and PKC α activity attenuates CFA-induced increases in GluA2 phosphorylation at Ser⁸⁸⁰ and cytosolic GluA2 expression in the dorsal horn^[23] (Fig. 1). Furthermore, blocking the CFA-induced increase in phosphorylation of GluA2 Ser⁸⁸⁰ reduces CFA-evoked GluA2 internalization in dorsal horn neurons^[23]. The increased intracellular Ca²⁺ concentration that results from Ca²⁺ influx through activated NMDA receptors may activate PKC α to phosphorylate GluA2 at Ser⁸⁸⁰. This phosphorylation would disrupt GluA2 bind-

ing to ABP/GRIP and promote GluA2 internalization in dorsal horn neurons. Indeed, the binding affinity of GluA2 to GRIP1 is significantly reduced in dorsal horn neurons after intraplantar CFA injection^[23]. Moreover, NMDA receptor-triggered GluA2 internalization requires Ca²⁺ influx directly through the NMDA receptor channel^[48,49]. AMPA complexes are physically coupled to NMDA receptor complexes by a PSD-95–stargazin linkage with GluA1/2/4 and NR2A/NR2B and to PKC α by PICK1 with GluA2 in the dorsal horn^[23]. This coupling may provide a molecular basis for an NMDA receptor–PKC α –AMPA signal cascade and enable Ca²⁺ to reach PKC α through NMDA receptor channel influx. The result would be phosphorylation of both GluA1 and GluA2 and changes in trafficking of both subunits. This conclusion is supported by the evidence that both GluA1 membrane insertion and GluA2 internalization require NMDA receptor activation in central neurons^[23,48-53]. Therefore, NMDA receptor/PKC α activation may be responsible for CFA-induced changes of GluA1 and GluA2 trafficking in dorsal horn neurons after inflammation.

It should be noted that the binding motif in the GluA2 C-terminus also binds other proteins that are involved in the regulation of synaptic GluA2 trafficking. For example, GluA2 interacts with N-ethylmaleimide-sensitive fusion (NSF) protein. This protein maintains the synaptic expression of GluA2-containing AMPARs, displaces PICK1 from the PICK1–GluA2 complex, and facilitates the delivery or stabilization of GluA2 at the plasma membrane^[54-56]. Interestingly, CFA-induced peripheral inflammation decreases NSF expression in the spinal cord^[24]. Whether and how this decrease contributes to CFA-induced dorsal horn GluA2 internalization is unknown and needs additional study.

5 Changes in dorsal horn AMPAR subunit trafficking participate in central sensitization during inflammatory pain

The evidence presented so far indicates that peripheral inflammatory insults change AMPAR subunit trafficking and functional properties at synaptic and extrasynaptic




Fig. 1. Protein kinase C alpha (PKC α)-dependent GluA2 internalization in dorsal horn 1 day after complete Freund's adjuvant (CFA) injection into rat hind-paw. Two intrathecal injections of the selective PKC α inhibitor Go6976 (Go, 25 μ g) given at a 12-h interval blocked the CFA-induced increase in GluA2 in the 150 k-g dorsal horn fraction 1 day post CFA injection (unpublished data). $n = 4$ /group. ** $P < 0.01$ vs V + S group; ## $P < 0.01$ vs V + C group. C, CFA; S, saline; V, vehicle (20% DMSO).

sites of dorsal horn neurons. Lastly, I will review the literature to address whether these changes are involved in the development and maintenance of inflammation-induced pain hypersensitivity. It has been reported that the induction of dorsal horn GluA1 membrane insertion by capsaicin or carrageenan correlates with the development of pain hypersensitivity^[19,21]. Similarly, the GluA2 internalization and GluA1 membrane insertion that occur 1 to 3 days after peripheral CFA injection correlate with the maintenance of CFA-induced pain hypersensitivity^[9,23,25]. Hence, inflammation-induced changes in dorsal horn AMPAR subunit trafficking are likely to participate in central sensitization during inflammatory pain. This conclusion is strongly supported by the following evidence. First, the activation of dorsal horn NMDA receptors and PKC α is required

for inflammation-induced dorsal horn GluA2 internalization during the maintenance of inflammatory pain (Fig. 1)^[4,23]; second, our recent work showed that dorsal horn PKC α knockdown not only blocks CFA-induced increases in extrasynaptic GluA1-containing and Ca²⁺-permeable AMPARs in dorsal horn neurons, but also attenuates CFA-induced mechanical allodynia 1 day after CFA injection; third, interfering with dorsal horn GluA2 internalization by blocking the GluA2 PKC α phosphorylation site, inhibiting spinal cord PKC α , or disrupting the recruitment of PKC α to GluA2, alleviates CFA-induced inflammatory pain during the maintenance period^[23,26] (Figs. 1 and 2); and fourth, peripheral CFA injection increases the number of synaptic and extrasynaptic Ca²⁺-permeable AMPARs in the dorsal horn during the maintenance of inflammatory pain^[23-25,28]. Given




Fig. 2. Protein kinase C alpha (PKC α)-dependent mechanical and thermal hypersensitivity 1 day after complete Freund's adjuvant (CFA) injection into rat hind-paw. Two intrathecal injections of Go6976 given at a 12-h interval dose-dependently attenuated CFA-induced mechanical allodynia (A) and thermal hyperalgesia (B) on the ipsilateral side. At the doses used, Go6976 did not affect basal paw withdrawal responses to mechanical (C) or thermal (D) stimuli on the contralateral side (unpublished data). $n = 5-6/\text{group}$. * $P < 0.05$, ** $P < 0.01$ vs corresponding baseline; # $P < 0.05$, ## $P < 0.01$ vs vehicle group.




Fig. 3. A proposed model for the mechanism by which changes in the trafficking of dorsal horn α -amino-3-hydroxy-5-methyl-4-isoxazolepropionic receptors (AMPA) contribute to inflammation-induced central sensitization. Under inflammatory conditions, dorsal horn N-methyl-D-aspartic acid receptor (NMDAR) activation increases Ca^{2+} influx through NMDAR channels. The elevation of intracellular Ca^{2+} initiates or potentiates intracellular Ca^{2+} -dependent cascades, including activation of protein kinases (e.g., PKC, PKA, or CaMKII). These cascades generate central sensitization as well as induce phosphorylation of AMPAR subunits GluA1 and GluA2 and promote GluA1 exocytosis and GluA2 endocytosis in dorsal horn neurons. The changes in AMPAR subunit composition result in an increase in AMPAR Ca^{2+} permeability in dorsal horn neurons. The increased permeability leads to more Ca^{2+} influx, which further enhances the activation of Ca^{2+} -dependent cascades. This positive feedback loop may be a key contributor to the maintenance of pain hypersensitivity under inflammatory pain conditions. PKA, protein kinase A; PKC α , protein kinase C alpha.

that the NMDA receptor, Ca^{2+} , and PKC are key players in inflammation-induced dorsal horn central sensitization^[4,57,58], I propose a model in which GluA1 membrane insertion and GluA2 internalization in the dorsal horn participate in the mechanism underlying the maintenance of central sensitization in inflammatory pain (Fig. 3). In brief, after inflammation, NMDA receptor activation induces Ca^{2+} influx in dorsal horn neurons. The increase in intracellular Ca^{2+} initiates and/or potentiates various Ca^{2+} -dependent intracellular cascades (including activation of protein kinases, e.g. PKC), induces phosphorylation of both GluA1 and GluA2, and promotes GluA1 membrane insertion and GluA2 internalization. The change in synaptic and extrasynaptic AMPAR subunit composition leads to increases in AMPAR Ca^{2+} permeability. The increased

permeability produces more Ca^{2+} influx, which further enhances the activation of Ca^{2+} -dependent intracellular cascades in the dorsal horn. This positive feedback loop that occurs through changes in AMPAR subunit trafficking and increases in the number of Ca^{2+} -permeable AMPARs in dorsal horn neurons may be critical for the maintenance of pain hypersensitivity in persistent inflammatory pain (Fig. 3).

6 Conclusion

Persistent (or chronic) pain is not simply the consequence of peripheral inputs or pathology but is also a reflection of changes in the plasticity of central nociceptive circuits (that is, central sensitization). These changes are sufficient to reduce the pain threshold, increase the magnitude and duration of responses to noxious input, and

permit normally innocuous inputs to induce pain sensation. Thus, central sensitization is a major contributor to many clinical pain syndromes. New research has revealed that profound changes occur in dorsal horn AMPAR subunit trafficking in persistent inflammatory pain. These trafficking changes contribute to the induction and/or maintenance of inflammation-induced central sensitization. The development of specific and selective drugs that interfere with changes in synaptic and extrasynaptic AMPAR trafficking may provide an effective novel strategy for preventing and/or treating persistent pain.

Acknowledgements: This work was supported by grants from the National Institutes of Health (NS058886 and NS072206), Rita Allen Foundation, Mr. David Koch and the Patrick C. Walsh Prostate Cancer Research Fund, and the Blaustein Pain Research Fund. The author thanks C.F. Levine, MS, ELS, for editorial assistance.

References:

- [1] Scholz J, Woolf CJ. Can we conquer pain? *Nat Neurosci* 2002, 5 Suppl: 1062–1067.
- [2] Burnashev N, Monyer H, Seeburg PH, Sakmann B. Divalent ion permeability of AMPA receptor channels is dominated by the edited form of a single subunit. *Neuron* 1992, 8(1): 189–198.
- [3] Hollmann M, Heinemann S. Cloned glutamate receptors. *Annu Rev Neurosci* 1994, 17: 31–108.
- [4] Tao YX. Dorsal horn alpha-amino-3-hydroxy-5-methyl-4-isoxazolepropionic acid receptor trafficking in inflammatory pain. *Anesthesiology* 2010, 112(5): 1259–1265.
- [5] Mao J, Price DD, Hayes RL, Lu J, Mayer DJ. Differential roles of NMDA and non-NMDA receptor activation in induction and maintenance of thermal hyperalgesia in rats with painful peripheral mononeuropathy. *Brain Res* 1992, 598(1–2): 271–278.
- [6] Garry EM, Moss A, Rosie R, Delaney A, Mitchell R, Fleetwood-Walker SM. Specific involvement in neuropathic pain of AMPA receptors and adapter proteins for the GluR2 subunit. *Mol Cell Neurosci* 2003, 24(1): 10–22.
- [7] Sorkin LS, Yaksh TL, Doom CM. Mechanical allodynia in rats is blocked by a Ca^{2+} permeable AMPA receptor antagonist. *Neuroreport* 1999, 10(17): 3523–3526.
- [8] Sorkin LS, Yaksh TL, Doom CM. Pain models display differential sensitivity to Ca^{2+} -permeable non-NMDA glutamate receptor antagonists. *Anesthesiology* 2001, 95(4): 965–973.
- [9] Park JS, Yaster M, Guan X, Xu JT, Shih MH, Guan Y, *et al.* Role of spinal cord alpha-amino-3-hydroxy-5-methyl-4-isoxazolepropionic acid receptors in complete Freund's adjuvant-induced inflammatory pain. *Mol Pain* 2008, 4(1): 67.
- [10] Seagrove LC, Suzuki R, Dickenson AH. Electrophysiological characterisations of rat lamina I dorsal horn neurones and the involvement of excitatory amino acid receptors. *Pain* 2004, 108(1–2): 76–87.
- [11] Stanfa LC, Hampton DW, Dickenson AH. Role of Ca^{2+} -permeable non-NMDA glutamate receptors in spinal nociceptive transmission. *Neuroreport* 2000, 11(14): 3199–3202.
- [12] Hartmann B, Ahmadi S, Heppenstall PA, Lewin GR, Schott C, Borchardt T, *et al.* The AMPA receptor subunits GluR-A and GluR-B reciprocally modulate spinal synaptic plasticity and inflammatory pain. *Neuron* 2004, 44(4): 637–650.
- [13] Collingridge GL, Isaac JT, Wang YT. Receptor trafficking and synaptic plasticity. *Nat Rev Neurosci* 2004, 5(12): 952–962.
- [14] Song I, Huganir RL. Regulation of AMPA receptors during synaptic plasticity. *Trends Neurosci* 2002, 25(11): 578–588.
- [15] Engelmann HS, Allen TB, MacDermott AB. The distribution of neurons expressing calcium-permeable AMPA receptors in the superficial laminae of the spinal cord dorsal horn. *J Neurosci* 1999, 19(6): 2081–2089.
- [16] Fletcher EJ, Lodge D. New developments in the molecular pharmacology of alpha-amino-3-hydroxy-5-methyl-4-isoxazole propionate and kainate receptors. *Pharmacol Ther* 1996, 70(1): 65–89.
- [17] Furuyama T, Kiyama H, Sato K, Park HT, Maeno H, Takagi H, *et al.* Region-specific expression of subunits of ionotropic glutamate receptors (AMPA-type, KA-type and NMDA receptors) in the rat spinal cord with special reference to nociception. *Brain Res Mol Brain Res* 1993, 18(1–2): 141–151.
- [18] Polgar E, Al Ghamdi KS, Todd AJ. Two populations of neurokinin 1 receptor-expressing projection neurons in lamina I of the rat spinal cord that differ in AMPA receptor subunit composition and density of excitatory synaptic input. *Neuroscience* 2010, 167(4): 1192–1204.
- [19] Galan A, Laird JM, Cervero F. *In vivo* recruitment by painful stimuli of AMPA receptor subunits to the plasma membrane of spinal cord neurons. *Pain* 2004, 112(3): 315–323.
- [20] Larsson M, Broman J. Translocation of GluR1-containing AMPA receptors to a spinal nociceptive synapse during acute noxious stimulation. *J Neurosci* 2008, 28(28): 7084–7090.
- [21] Choi JI, Svensson CI, Koehn FJ, Bhushute A, Sorkin LS. Peripheral inflammation induces tumor necrosis factor dependent AMPA receptor trafficking and Akt phosphorylation in spinal cord in addition to pain behavior. *Pain* 2010, 149(2): 243–253.
- [22] Pezet S, Marchand F, D'Mello R, Grist J, Clark AK, Malcangio M, *et al.* Phosphatidylinositol 3-kinase is a key mediator of central

- sensitization in painful inflammatory conditions. *J Neurosci* 2008, 28(16): 4261–4270.
- [23] Park JS, Voitenko N, Petralia RS, Guan X, Xu JT, Steinberg JP, *et al.* Persistent inflammation induces GluR2 internalization via NMDA receptor-triggered PKC activation in dorsal horn neurons. *J Neurosci* 2009, 29(10): 3206–3219.
- [24] Katano T, Furue H, Okuda-Ashitaka E, Tagaya M, Watanabe M, Yoshimura M, *et al.* N-ethylmaleimide-sensitive fusion protein (NSF) is involved in central sensitization in the spinal cord through GluR2 subunit composition switch after inflammation. *Eur J Neurosci* 2008, 27(12): 3161–3170.
- [25] Kopach O, Kao SC, Petralia RS, Belan P, Tao YX, Voitenko N. Inflammation alters trafficking of extrasynaptic AMPA receptors in tonically firing lamina II neurons of the rat spinal dorsal horn. *Pain* 2011, 152(4): 912–923.
- [26] Atianjoh FE, Yaster M, Zhao X, Takamiya K, Xia J, Gauda EB, *et al.* Spinal cord protein interacting with C kinase 1 is required for the maintenance of complete Freund's adjuvant-induced inflammatory pain but not for incision-induced post-operative pain. *Pain* 2010, 151(1): 226–234.
- [27] Wang W, Petralia RS, Takamiya K, Xia J, Li YQ, Huganir RL, *et al.* Preserved acute pain and impaired neuropathic pain in mice lacking protein interacting with C Kinase 1. *Mol Pain* 2011, 7: 11.
- [28] Vikman KS, Rycroft BK, Christie MJ. Switch to Ca^{2+} -permeable AMPA and reduced NR2B NMDA receptor-mediated neurotransmission at dorsal horn nociceptive synapses during inflammatory pain in the rat. *J Physiol* 2008, 586(2): 515–527.
- [29] Banke TG, Bowie D, Lee H, Huganir RL, Schousboe A, Traynelis SF. Control of GluR1 AMPA receptor function by cAMP-dependent protein kinase. *J Neurosci* 2000, 20(1): 89–102.
- [30] Man HY, Sekine-Aizawa Y, Huganir RL. Regulation of {alpha}-amino-3-hydroxy-5-methyl-4-isoxazolepropionic acid receptor trafficking through PKA phosphorylation of the Glu receptor 1 subunit. *Proc Natl Acad Sci U S A* 2007, 104(9): 3579–3584.
- [31] Oh MC, Derkach VA, Guire ES, Soderling TR. Extrasynaptic membrane trafficking regulated by GluR1 serine 845 phosphorylation primes AMPA receptors for long-term potentiation. *J Biol Chem* 2006, 281(2): 752–758.
- [32] Derkach V, Barria A, Soderling TR. Ca^{2+} /calmodulin-kinase II enhances channel conductance of alpha-amino-3-hydroxy-5-methyl-4-isoxazolepropionate type glutamate receptors. *Proc Natl Acad Sci U S A* 1999, 96(6): 3269–3274.
- [33] Hayashi Y, Shi SH, Esteban JA, Piccini A, Ponce JC, Malinow R. Driving AMPA receptors into synapses by LTP and CaMKII: requirement for GluR1 and PDZ domain interaction. *Science* 2000, 287(5461): 2262–2267.
- [34] Boehm J, Kang MG, Johnson RC, Esteban J, Huganir RL, Malinow R. Synaptic incorporation of AMPA receptors during LTP is controlled by a PKC phosphorylation site on GluR1. *Neuron* 2006, 51(2): 213–225.
- [35] Shen L, Liang F, Walensky LD, Huganir RL. Regulation of AMPA receptor GluR1 subunit surface expression by a 4.1N-linked actin cytoskeletal association. *J Neurosci* 2000, 20(21): 7932–7940.
- [36] Lin DT, Makino Y, Sharma K, Hayashi T, Neve R, Takamiya K, *et al.* Regulation of AMPA receptor extrasynaptic insertion by 4.1N, phosphorylation and palmitoylation. *Nat Neurosci* 2009, 12(7): 879–887.
- [37] Fang L, Wu J, Zhang X, Lin Q, Willis WD. Increased phosphorylation of the GluR1 subunit of spinal cord alpha-amino-3-hydroxy-5-methyl-4-isoxazole propionate receptor in rats following intradermal injection of capsaicin. *Neuroscience* 2003, 122(1): 237–245.
- [38] Fang L, Wu J, Lin Q, Willis WD. Protein kinases regulate the phosphorylation of the GluR1 subunit of AMPA receptors of spinal cord in rats following noxious stimulation. *Brain Res Mol Brain Res* 2003, 118(1–2): 160–165.
- [39] Lu Y, Sun YN, Wu X, Sun Q, Liu FY, Xing GG, *et al.* Role of alpha-amino-3-hydroxy-5-methyl-4-isoxazolepropionate (AMPA) receptor subunit GluR1 in spinal dorsal horn in inflammatory nociception and neuropathic nociception in rat. *Brain Res* 2008, 1200: 19–26.
- [40] Dong H, O'Brien RJ, Fung ET, Lanahan AA, Worley PF, Huganir RL. GRIP: a synaptic PDZ domain-containing protein that interacts with AMPA receptors. *Nature* 1997, 386(6622): 279–284.
- [41] Dong H, Zhang P, Song I, Petralia RS, Liao D, Huganir RL. Characterization of the glutamate receptor-interacting proteins GRIP1 and GRIP2. *J Neurosci* 1999, 19(16): 6930–6941.
- [42] Dong H, Zhang P, Liao D, Huganir RL. Characterization, expression, and distribution of GRIP protein. *Ann N Y Acad Sci* 1999, 868: 535–540.
- [43] Dev KK, Nakanishi S, Henley JM. The PDZ domain of PICK1 differentially accepts protein kinase C-alpha and GluR2 as interacting ligands. *J Biol Chem* 2004, 279(40): 41393–41397.
- [44] Perez JL, Khatri L, Chang C, Srivastava S, Osten P, Ziff EB. PICK1 targets activated protein kinase Calpha to AMPA receptor clusters in spines of hippocampal neurons and reduces surface levels of the AMPA-type glutamate receptor subunit 2. *J Neurosci* 2001, 21(15): 5417–5428.
- [45] Staudinger J, Lu J, Olson EN. Specific interaction of the PDZ domain protein PICK1 with the COOH terminus of protein kinase C-alpha. *J Biol Chem* 1997, 272(51): 32019–32024.
- [46] Chung HJ, Xia J, Scannevin RH, Zhang X, Huganir RL. Phosphorylation of the AMPA receptor subunit GluR2 differentially regulates its interaction with PDZ domain-containing proteins. *J Neurosci* 2000, 20(19): 7258–7267.
- [47] Matsuda S, Mikawa S, Hirai H. Phosphorylation of serine-880 in GluR2 by protein kinase C prevents its C terminus from binding

- with glutamate receptor-interacting protein. *J Neurochem* 1999, 73(4): 1765–1768.
- [48] Iwakura Y, Nagano T, Kawamura M, Horikawa H, Ibaraki K, Takei N, *et al.* N-methyl-D-aspartate-induced alpha-amino-3-hydroxy-5-methyl-4-isoxazolepropionic acid (AMPA) receptor down-regulation involves interaction of the carboxyl terminus of GluR2/3 with Pick1. Ligand-binding studies using Sindbis vectors carrying AMPA receptor decoys. *J Biol Chem* 2001, 276(43): 40025–40032.
- [49] Tigaret CM, Thalhammer A, Rast GF, Specht CG, Auberson YP, Stewart MG, *et al.* Subunit dependencies of N-methyl-D-aspartate (NMDA) receptor-induced alpha-amino-3-hydroxy-5-methyl-4-isoxazolepropionic acid (AMPA) receptor internalization. *Mol Pharmacol* 2006, 69(4): 1251–1259.
- [50] Moulton PR, Cross A, Santos SD, Carvalho AL, Lindsay Y, Connolly CN, *et al.* Leptin regulates AMPA receptor trafficking via PTEN inhibition. *J Neurosci* 2010, 30(11): 4088–4101.
- [51] Corera AT, Doucet G, Fon EA. Long-term potentiation in isolated dendritic spines. *PLoS One* 2009, 4(6): e6021.
- [52] Serulle Y, Zhang S, Ninan I, Puzzo D, McCarthy M, Khatri L, *et al.* A GluR1-cGKII interaction regulates AMPA receptor trafficking. *Neuron* 2007, 56(4): 670–688.
- [53] Kim MJ, Dunah AW, Wang YT, Sheng M. Differential roles of NR2A- and NR2B-containing NMDA receptors in Ras-ERK signaling and AMPA receptor trafficking. *Neuron* 2005, 46(5): 745–760.
- [54] Braithwaite SP, Xia H, Malenka RC. Differential roles for NSF and GRIP/ABP in AMPA receptor cycling. *Proc Natl Acad Sci U S A* 2002, 99(10): 7096–7101.
- [55] Hanley JG, Khatri L, Hanson PI, Ziff EB. NSF ATPase and alpha-/beta-SNAPs disassemble the AMPA receptor-PICK1 complex. *Neuron* 2002, 34(1): 53–67.
- [56] Huang Y, Man HY, Sekine-Aizawa Y, Han Y, Juluri K, Luo H, *et al.* S-nitrosylation of N-ethylmaleimide sensitive factor mediates surface expression of AMPA receptors. *Neuron* 2005, 46(4): 533–540.
- [57] Ji RR, Kohno T, Moore KA, Woolf CJ. Central sensitization and LTP: do pain and memory share similar mechanisms? *Trends Neurosci* 2003, 26(12): 696–705.
- [58] Latremoliere A, Woolf CJ. Central sensitization: a generator of pain hypersensitivity by central neural plasticity. *J Pain* 2009, 10(9): 895–926.