

Prefrontal cortical α_{2A} -adrenoceptors and a possible primate model of attention deficit and hyperactivity disorder

Chao-Lin Ma, Xuan Sun, Fei Luo, Bao-Ming Li

Center for Neuropsychiatric Disorders, Institute of Life Science, Nanchang University, Nanchang 330031, China

Corresponding author: Chao-Lin Ma. E-mail: chaolin.ma@gmail.com

© Shanghai Institutes for Biological Sciences, CAS and Springer-Verlag Berlin Heidelberg 2015

Attention deficit and hyperactivity disorder (ADHD), a prevalent syndrome in children worldwide, is characterized by impulsivity, inappropriate inattention, and/or hyperactivity. It seriously afflicts cognitive development in childhood, and may lead to chronic under-achievement, academic failure, problematic peer relationships, and low self-esteem. There are at least three challenges for the treatment of ADHD. First, the neurobiological bases of its symptoms are still not clear. Second, the commonly prescribed medications, most showing short-term therapeutic efficacy but with a high risk of serious side-effects, are mainly based on a dopamine mechanism. Third, more novel and efficient animal models, especially in nonhuman primates, are required to accelerate the development of new medications. In this article, we review research progress in the related fields, focusing on our previous studies showing that blockade of prefrontal cortical α_{2A} -adrenoceptors in monkeys produces almost all the typical behavioral symptoms of ADHD.

Keywords: prefrontal cortex; α_{2A} -adrenoceptors; cognitive functions; attention deficit and hyperactivity disorder; animal models

Introduction

Attention deficit and hyperactivity disorder (ADHD) is one of the most prevalent childhood neurodevelopmental conditions, affecting 3–5% of grade-school children worldwide^[1]. It is characterized by inappropriate levels of inattention, impulsivity, and/or hyperactivity^[2–4]. These symptoms develop in childhood, and can persist into adolescence and adulthood^[5]. ADHD seriously affects cognitive development^[6–8], and, without appropriate treatment, has consequences for the risk of anxiety, substance abuse, and depression in adulthood^[2, 5, 9].

The neurobiological bases of ADHD symptoms are still not clear^[10]. Clarifying them can help better understand the biological vulnerabilities that may underlie ADHD in a specific patient and how to modulate the responses to treatment, thereby contributing to better and more effective therapy.

It has been suggested that the symptoms involve a dopaminergic mechanism in the prefrontal cortex (PFC)

and striatum^[11, 12]. Experimentally decreased dopamine (DA) release in the PFC results in ADHD-like symptoms^[13, 14]. To date, DA dysregulation is thought to be central to the neurobiology of ADHD, and its pharmacological treatment, such as methylphenidate (MPH, i.e. Ritalin)^[15–17], levels the DA concentration in the synapse and extrasynaptic space in the PFC as a blocker of the DA transporter. MPH ameliorates inappropriate inattention^[18–20], decreases impulsivity^[21], and enhances inhibitory control^[22]. However, as MPH is a prescription psychostimulant, there are strong concerns over drug dependence, paranoia, schizophrenia, and behavioral sensitization that might be caused by long-term therapy, similar to other stimulants^[23–25].

Converging evidence indicates that the pathophysiology of ADHD has multiple origins^[26–32]; for instance, norepinephrine (NE) has long been implicated^[33, 34]. In this paper, we review research progress in the relevant fields, focusing on the potential relationship between prefrontal α_{2A} -adrenoceptors and ADHD in nonhuman primates^[35–39].

Prefrontal Cognitive Dysfunctions in ADHD

The PFC plays a key role in cognitive functions such as working memory, the regulation of attention, and behavioral inhibition. Imaging and neuropsychological studies have shown that patients with ADHD have poor PFC functions, including poor attention regulation^[40], limited working memory^[41], and inability to inhibit inappropriate motor activity^[42].

Working memory is a fundamental higher-order function, underlies a wide range of executive functional processes^[43, 44], and is primarily controlled by the PFC^[45, 46]. It has been shown that ADHD patients have altered architecture and less activation in the PFC^[47–49]. Persistent working-memory problems are the main cognitive deficit in ADHD^[40, 41, 50, 51].

Attention brings sensory or mental stimuli to the forefront of awareness^[40, 52], and plays a pivotal role in mediating the executive functions of the PFC. During distracted states, the capacity to diminish the awareness of relevant stimuli is compromised. Compared to normal peers, ADHD patients show attention deficits in detecting invalidly-cued targets with slower speed and less accuracy^[53, 54].

Inhibitory control of behavior is one of the most important functions of the PFC^[55]; ablation or lesion of the frontal cortex in monkeys induces locomotor hyperactivity^[56–58]. Perhaps the most fundamental deficit in ADHD is the lack of response inhibition^[52]. In laboratory studies of tasks that measure inhibitory control, children with ADHD often perform more poorly than both normal controls and children with other psychiatric disorders^[59, 60]. Schulz *et al.* reported that response inhibition in adolescents diagnosed with ADHD is primarily mediated by fronto-striatal circuitry^[61, 62].

Prefrontal α_{2A} -Adrenoceptor Blockade Produces ADHD Phenotypes

ADHD has been posited to be caused by hypofunctional catecholamine systems^[63] in multiple brain regions including the PFC^[64–66] and striatum^[67]. Implicated in this are NE projections that originate primarily from neurons in the locus coeruleus and send projections to multiple regions, including the PFC^[68]. There are many subtypes of adrenergic receptors in the PFC, including the α_{2A} subtype.

The α_{2A} -adrenoceptors are localized at both pre- and post-synaptic NE terminals^[69]. However, studies in rodents, monkeys, and humans have shown that lower to moderate levels of NE have a beneficial influence on prefrontal cognitive functions through action at post-synaptic α_{2A} -adrenoceptors^[64, 70].

ADHD symptoms can be mimicked by blockade of α_{2A} -adrenoceptors in the PFC. To investigate the role of prefrontal α_{2A} -adrenoceptors in the inhibitory control of behavior, we trained two monkeys to perform a go/no-go task, and the α_{2A} -adrenergic antagonist yohimbine was infused bilaterally and chronically into the dorsolateral PFC with mini-osmotic pumps. We found that blockade of the α_{2A} -adrenoceptors selectively impaired the “no-go” performance of monkeys, leaving the “go” performance intact. In quite a few cases, the monkeys should have kept their hands still and not touch the screen (no-go), but they made a response to the screen^[38]. Infusion of saline at the same cortical locations did not affect the no-go performance, indicating that the yohimbine-induced impulse was not because of nonspecific factors such as infusion-induced cortical damage (Fig. 1A). Our previous work provided the first behavioral evidence that α_{2A} -adrenoceptors in the dorsolateral PFC are involved in the inhibitory control of behavior.

In addition, the monkeys' locomotor activity was monitored before, during and after yohimbine infusion into the dorsolateral PFC. Compared to that before administration, the daily locomotor activity increased dramatically during the 8-day administration of yohimbine; this gradually returned to normal after the infusion was stopped (Fig. 1B). Infusion of saline at the same location did not cause locomotor hyperactivity^[39]. This work suggests that the α_{2A} -adrenoceptors in the dorsolateral PFC are associated with locomotor activity, and the dorsolateral PFC dysfunction of α_{2A} -adrenergic transmission could be one of the main causes of the impulsive behaviors and hyperactivity in children with ADHD.

Due to the limitations of working on nonhuman primates, we also implemented similar experiments on rats to assess the dose-dependent and age effects of yohimbine at a homologous cortical site, the medial PFC. The results showed that yohimbine infused into the medial PFC dose-dependently induced hyperactivity in rats of different ages, and the trends showed that the younger the rats, the more hyperactivity




Fig. 1. Yohimbine infused bilaterally and chronically into the dorsolateral prefrontal cortex impairs impulse control and induces locomotor hyperactivity (adapted from Ma CL *et al.*, *Neuroreport*, 2003^[38] and *Biol Psychiatry*, 2005^[39]). (A) Yohimbine impairs “no-go” performance but has no effect on “go” performance. In several cases, the monkeys should not touch the screen (no-go), but they make a response. (B) Daily locomotor activity increases during administration of yohimbine. Each trace is a daily recording from 06:00 to 18:00. Inset: Reconstructed sites for chronic administration of yohimbine and saline. Filled symbols, yohimbine infusion; open symbols, saline infusion; as, arcuate sulcus; ps, principal sulcus.

presented at the same dose (unpublished data). All these results showed that dysfunction of the PFC α_{2A} -adrenoceptors results in the behavioral problems seen in ADHD.

ADHD symptoms can also be induced in humans by reducing the stimulation of α_{2A} -adrenoceptors. Kopeckova *et al.* investigated a polymorphism in the promoter region of the gene encoding DA beta-hydroxylase, an enzyme that reduces NE synthesis, and found that the affected children had poor sustained attention, weaker impulse control, and impaired executive function^[71]. Genetic alterations in α_{2A} -adrenoceptors also impair PFC executive function, and lead to conditions seen in ADHD^[72]. Thus, prefrontal α_{2A} -adrenoceptors are required for attention and behaviors in humans too.

Prefrontal α_{2A} -Adrenoceptor Stimulation Ameliorates Cognitive Dysfunctions in ADHD

Behavioral, pharmacological, and electrophysiological

research has shown that stimulation of α_{2A} -adrenoceptors has a beneficial influence on PFC cognitive functions. Arnsten *et al.* found that systemic administration of the α_{2A} -adrenergic agonist guanfacine improves working memory in monkeys^[73]. Steere demonstrated that systemic administration of guanfacine improves visual object discrimination reversal performance in aged rhesus monkeys^[74]. Our work showed that both systemic administration and local infusion of guanfacine into the PFC improve visuomotor associative learning^[70, 75]. Using an iontophoretic technique, stimulation of α_{2A} -adrenoceptors in the PFC was found to increase the spiking activity associated with working memory in behaving monkeys^[37, 76].

Neuronal activity in the PFC associated with working memory can be enhanced by α_{2A} -adrenoceptor stimulation through cAMP-HCN signaling pathways^[76, 77]. Our work suggested that under normal physiological conditions, the α_{2A} -adrenoceptors in pyramidal cells can be activated through Gi-cAMP-HCN signaling^[78]. On the other hand,

under stress, activation of α_{2A} -adrenoceptors to protect PFC functions might occur *via* the Gi-cAMP-PKA-CaMKII-AMPA signaling pathway^[76]. Both mechanisms together optimize the synaptic inputs to pyramidal neurons and determine the synaptic outputs for PFC cognitive functions. Indeed, it has been reported that guanfacine at relatively high doses suppresses evoked excitatory postsynaptic currents, and has no enhanced effect or even suppresses delay-related activity^[76].

New Insights to Develop a Primate Model of ADHD

As noted above, most of the commonly-prescribed medications for ADHD are psychostimulants, which are reported to have short-term therapeutic efficacy but with a high risk of serious adverse effects with long-term treatment. It is urgent to find new medications with high therapeutic efficacy and low adverse effects for children with ADHD. The first step now is to develop novel animal models of ADHD. A good model should very nearly show the fundamental behavioral characteristics of ADHD, conform to a theoretical rationale for ADHD, account for the neurobiology, and respond to therapeutic interventions both behaviorally and pharmacologically^[79].

Currently, animal models of ADHD are genetic and non-genetic^[80]. The spontaneously-hypertensive rat (SHR), the most widely used model, is a genetic model^[81, 82]. SHRs exhibit hyperactivity^[83, 84], impulsivity/inattention^[82, 85], and poor learning and memory^[86]. They also have disturbances in glutamate, DA, and NE functions, which in parallel demonstrate that ADHD patients have defects in the neuronal circuits required for reward-guided associative learning and memory formation^[87]. Clearly, the SHR is a good model for the study of memory deficits in ADHD, primarily in the context of particular risk factors/symptoms, responsiveness to specific drugs or other treatments or biomarkers for the diagnosis of ADHD, and for understanding the pathological mechanisms for the development of therapeutic approaches. However, SHRs do not fulfill all the behavioral and pharmacological profiles of an ADHD model; for example, ADHD-like behaviors in SHRs are not restricted to males^[88]. Hyperactive behavior in SHRs is ameliorated only by high doses of amphetamine or MPH^[84], unlike ADHD patients, whose behavioral deficits

can be improved with low doses of MPH. Importantly, ADHD patients show reduced regional cerebral blood flow in the frontal cortex^[89], while SHRs do not^[90, 91].

Our previous research with monkeys indicates that blockade of the prefrontal α_{2A} receptors induces locomotor hyperactivity, impulsivity, and poor attention regulation/working memory. These results verify the feasibility and acceptability of treating ADHD by stimulating α_{2A} -adrenoceptors in the PFC or up-regulating the NE concentration in synapses and extrasynaptic space in the PFC. Actually, the α_{2A} -adrenergic agonists guanfacine and clonidine have been used experimentally and clinically to treat ADHD^[92–97]. The selective inhibitor of the NE transporter atomoxetine (tomoxetine or LY139603) has also been reported to alleviate ADHD symptoms^[98]. All these medications have achieved much better therapeutic efficacy with less adverse effects^[99] than MPH and amphetamines, although there is controversy regarding the long-term effectiveness^[100, 101].

Thus, in the future, studies should focus on developing a novel ADHD model in nonhuman primates, by down-regulating or blocking the α_{2A} -adrenoceptors in the dorsolateral PFC. This could be realized by chronic bilateral infusion of yohimbine. This kind of animal model could approximate the fundamental behavioral characteristics of ADHD, conform to a theoretical rationale for ADHD associated with prefrontal α_{2A} -adrenoceptors, and account for the neurobiology and therapeutic interventions in terms of both pharmacological and behavioral functions.

ACKNOWLEDGMENTS

This review was supported by grants from the National Natural Science Foundation of China (31460263, 30990263, and 31271171), and the National Basic Research Development Program (973 Program) of China (2013CB835101).

Received date: 2014-06-30; Accepted date: 2015-02-11

REFERENCES

- [1] Graetz BW, Sawyer MG, Hazell PL, Arney F, Baghurst P. Validity of DSM-IVADHD subtypes in a nationally representative sample of Australian children and adolescents. *J Am Acad Child Adolesc Psychiatry* 2001, 40: 1410–1417.
- [2] Mannuzza S, Klein RG, Bessler A, Malloy P, Lapadula M. Adult outcome of hyperactive boys - educational-

- achievement, occupational rank, and psychiatric status. *Arch Gen Psychiatry* 1993, 50: 565–576.
- [3] Steinhoff KW. Special issues in the diagnosis and treatment of ADHD in adolescents. *Postgrad Med* 2008, 120: 60–68.
- [4] Biederman J. Attention-deficit/hyperactivity disorder: a selective overview. *Biol Psychiatry* 2005, 57: 1215–1220.
- [5] Kessler RC, Adler L, Barkley R, Biederman J, Conners CK, Demler O, *et al.* The prevalence and correlates of adult ADHD in the United States: results from the National Comorbidity Survey Replication. *Am J Psychiatry* 2006, 163: 716–723.
- [6] Huang-Pollock CL, Nigg JT, Carr TH. Deficient attention is hard to find: applying the perceptual load model of selective attention to attention deficit hyperactivity disorder subtypes. *J Child Psychol Psychiatry* 2005, 46: 1211–1218.
- [7] Waldie KE, Hausmann M. Right fronto-parietal dysfunction in children with ADHD and developmental dyslexia as determined by line bisection judgements. *Neuropsychologia* 2010, 48: 3650–3656.
- [8] ter Huurne N, Onnink M, Kan C, Franke B, Buitelaar J, Jensen O. Behavioral consequences of aberrant alpha lateralization in attention-deficit/hyperactivity disorder. *Biol Psychiatry* 2013, 74: 227–233.
- [9] Wilens TE, Biederman J, Spencer TJ. Attention deficit/hyperactivity disorder across the lifespan. *Annu Rev Med* 2002, 53: 113–131.
- [10] Li JJ, Lee SS. Negative emotionality mediates the association of 5-HTTLPR genotype and depression in children with and without ADHD. *Psychiatry Res* 2014, 215: 163–169.
- [11] Levy F, Swanson JM. Timing, space and ADHD: the dopamine theory revisited. *Aust N Z J Psychiatry* 2001, 35: 504–511.
- [12] Del Campo N, Chamberlain SR, Sahakian BJ, Robbins TW. The roles of dopamine and noradrenaline in the pathophysiology and treatment of attention-deficit/hyperactivity disorder. *Biol Psychiatry* 2011, 69: e145–157.
- [13] Bubser M SW. 6-Hydroxydopamine lesion of the rat prefrontal cortex increases locomotor activity, impairs acquisition of delayed alternation tasks, but does not affect uninterrupted tasks in the radial maze. *Behav Brain Res* 1990, 37: 157–168.
- [14] Arnsten AF. Stimulants: Therapeutic actions in ADHD. *Neuropsychopharmacology* 2006, 31: 2376–2383.
- [15] Greenhill LL, Halperin JM, Abikoff H. Stimulant medications. *J Am Acad Child Adolesc Psychiatry* 1999, 38: 503–512.
- [16] Safer DJ, Malever M. Stimulant treatment in Maryland public schools. *Pediatrics* 2000, 106: 533–539.
- [17] Swanson J, Baler RD, Volkow ND. Understanding the effects of stimulant medications on cognition in individuals with attention-deficit hyperactivity disorder: a decade of progress. *Neuropsychopharmacology* 2011, 36: 207–226.
- [18] Bellgrove MA, Barry E, Johnson KA, Cox M, Daibhis A, Daly M, *et al.* Spatial attentional bias as a marker of genetic risk, symptom severity, and stimulant response in ADHD. *Neuropsychopharmacology* 2008, 33: 2536–2545.
- [19] Nigg JT, Swanson JM, Hinshaw SP. Covert visual spatial attention in boys with attention deficit hyperactivity disorder: lateral effects, methylphenidate response and results for parents. *Neuropsychologia* 1997, 35: 165–176.
- [20] Sheppard DM, Bradshaw JL, Mattingley JB, Lee P. Effects of stimulant medication on the lateralisation of line bisection judgements of children with attention deficit hyperactivity disorder. *J Neurol Neurosurg Psychiatry* 1999, 66: 57–63.
- [21] Fallu A, Richard C., Prinzo R, Binder, C. Does OROS-methylphenidate improve core symptoms and deficits in executive function? Results of an open-label trial in adults with attention deficit hyperactivity disorder. *Curr Med Res Opin* 2006, 22: 2557–2566.
- [22] Mehta MA, Owen AM, Sahakian BJ, Mavaddat N, Pickard JD, Robbins TW. Methylphenidate enhances working memory by modulating discrete frontal and parietal lobe regions in the human brain. *J Neurosci* 2000, 20: RC65.
- [23] Kociancic T, Reed MD, Findling RL. Evaluation of risks associated with short- and long-term psychostimulant therapy for treatment of ADHD in children. *Expert Opin Drug Saf* 2004, 3: 93–100.
- [24] Pappadopulos E, Woolston S, Chait A, Perkins M, Connor DF, Jensen PS. Pharmacotherapy of aggression in children and adolescents: efficacy and effect size. *J Can Acad Child Adolesc Psychiatry* 2006, 15: 27–39.
- [25] Brown RT, Amler RW, Freeman WS, Perrin JM, Stein MT, Feldman HM, *et al.* Treatment of attention-deficit/hyperactivity disorder: overview of the evidence. *Pediatrics* 2005, 115: e749–757.
- [26] Sharma A, Couture J. A review of the pathophysiology, etiology, and treatment of attention-deficit hyperactivity disorder (ADHD). *Ann Pharmacother* 2014, 48: 209–225.
- [27] Fusar-Poli P, Rubia K, Rossi G, Sartori G, Balottin U. Striatal dopamine transporter alterations in ADHD: pathophysiology or adaptation to psychostimulants? A meta-analysis. *Am J Psychiatry* 2012, 169: 264–272.
- [28] Bock J, Braun K. The impact of perinatal stress on the functional maturation of prefronto-cortical synaptic circuits: implications for the pathophysiology of ADHD? *Prog Brain Res* 2011, 189: 155–169.
- [29] Pasini A, D'Agati E. Pathophysiology of NSS in ADHD. *World J Biol Psychiatry* 2009, 10: 495–502.
- [30] Soros P. Tourette syndrome, ADHD, and the limbic system: investigating the pathophysiology. *Dev Med Child Neurol* 2008, 50: 486.

- [31] Donnelly CL. History and pathophysiology of ADHD. *CNS Spectr* 2006, 11: 4–6.
- [32] Jensen PS. ADHD: current concepts on etiology, pathophysiology, and neurobiology. *Child Adolesc Psychiatr Clin N Am* 2000, 9: 557–572, vii–viii.
- [33] Zametkin AJ, Rapoport JL. Neurobiology of attention deficit disorder with hyperactivity: where have we come in 50 years? *J Am Acad Child Adolesc Psychiatry* 1987, 26: 676–686.
- [34] Arnsten AF, Steere JC, Hunt RD. The contribution of alpha 2-noradrenergic mechanisms of prefrontal cortical cognitive function. Potential significance for attention-deficit hyperactivity disorder. *Arch Gen Psychiatry* 1996, 53: 448–455.
- [35] Li BM, Mei ZT. Delayed-response deficit induced by local injection of the alpha-2 adrenergic antagonist yohimbine into the dorsolateral prefrontal cortex in young adult monkeys. *Behav Neural Biol* 1994, 62: 134–139.
- [36] Li BM, Kubota K. Alpha-2 adrenergic modulation of prefrontal cortical neuronal activity related to a visual discrimination task with GO and NO-GO performances in monkeys. *Neurosci Res* 1998, 31: 83–95.
- [37] Li BM, Mao ZM, Wang M, Mei ZT. Alpha-2 adrenergic modulation of prefrontal cortical neuronal activity related to spatial working memory in monkeys. *Neuropsychopharmacology* 1999, 21: 601–610.
- [38] Ma CL, Qi XL, Peng JY, Li BM. Selective deficit in no-go performance induced by blockade of prefrontal cortical alpha 2-adrenoceptors in monkeys. *Neuroreport* 2003, 14: 1013–1016.
- [39] Ma CL, Arnsten AFT, Li BM. Locomotor hyperactivity induced by blockade of prefrontal cortical alpha(2)-adrenoceptors in monkeys. *Biol Psychiatry* 2005, 57: 192–195.
- [40] Barkley RA. Behavioral inhibition, sustained attention, and executive functions: Constructing a unifying theory of ADHD. *Psychol Bull* 1997, 121: 65–94.
- [41] Martinussen R, Hayden J, Hogg-Johnson S, Tannock R. A meta-analysis of working memory impairments in children with attention-deficit/hyperactivity disorder. *J Am Acad Child Adolesc Psychiatry* 2005, 44: 377–384.
- [42] Oosterlaan J, Logan GD, Sergeant JA. Response inhibition in ADHD, CD, comorbid ADHD+CD, anxious, and control children: a meta-analysis of studies with the stop task. *J Child Psychol Psychiatry* 1998, 39: 411–425.
- [43] Roberts RJ, Pennington BF. An interactive framework for examining prefrontal cognitive processes. *Dev Neuropsychol* 1996, 12: 105–126.
- [44] Klingberg T, Fernell E, Olesen PJ, Johnson M, Gustafsson P, Dahlstrom K, *et al.* Computerized training of working memory in children with ADHD - A randomized, controlled trial. *J Am Acad Child Adolesc Psychiatry* 2005, 44: 177–186.
- [45] D'Esposito M, Detre JA, Alsop DC, Shin RK, Atlas S, Grossman M. The neural basis of the central executive system of working memory. *Nature* 1995, 378: 279–281.
- [46] Smith EE, Jonides J, Marshuetz C, Koeppel RA. Components of verbal working memory: evidence from neuroimaging. *Proc Natl Acad Sci U S A* 1998, 95: 876–882.
- [47] Rubia K, Overmeyer S, Taylor E, Brammer M, Williams SC, Simmons A, *et al.* Hypofrontality in attention deficit hyperactivity disorder during higher-order motor control: a study with functional MRI. *Am J Psychiatry* 1999, 156: 891–896.
- [48] Yeo RA, Hill DE, Campbell RA, Vigil J, Petropoulos H, Hart B, *et al.* Proton magnetic resonance spectroscopy investigation of the right frontal lobe in children with attention-deficit/hyperactivity disorder. *J Am Acad Child Adolesc Psychiatry* 2003, 42: 303–310.
- [49] Dickstein SG, Bannon K, Castellanos FX, Milham MP. The neural correlates of attention-deficit/hyperactivity disorder: an ALE metaanalysis. *J Child Psychol Psychiatry* 2006, 47: 1051–1062.
- [50] Rapport MD, Scanlan SW, Denney CB. Attention-deficit/hyperactivity disorder and scholastic achievement: a model of dual developmental pathways. *J Child Psychol Psychiatry* 1999, 40: 1169–1183.
- [51] Biederman J, Monuteaux MC, Doyle AE, Seidman LJ, Wilens TE, Ferrero F, *et al.* Impact of executive function deficits and attention-deficit/hyperactivity disorder (ADHD) on academic outcomes in children. *J Consult Clin Psychol* 2004, 72: 757–766.
- [52] Barkley RA, Grodzinsky G, DuPaul GJ. Frontal lobe functions in attention deficit disorder with and without hyperactivity: A review and research report. *J Abnorm Child Psychol* 1992, 20: 163–188.
- [53] Bellgrove MA, Eramudugolla R, Newman DP, Vance A, Mattingley JB. Influence of attentional load on spatial attention in acquired and developmental disorders of attention. *Neuropsychologia* 2013, 51: 1085–1093.
- [54] Bellgrove MA, Johnson KA, Barry E, Mulligan A, Hawi Z, Gill M, *et al.* Dopaminergic haplotype as a predictor of spatial inattention in children with attention-deficit/hyperactivity disorder. *Arch Gen Psychiatry* 2009, 66: 1135–1142.
- [55] Iversen SD, Mishkin M. Perseverative interference in monkeys following selective lesions of the inferior prefrontal convexity. *Exp Brain Res* 1970, 11: 376–386.
- [56] French GM. Locomotor effects of regional ablations of frontal cortex in rhesus monkeys. *J Comp Physiol Psychol* 1959, 52: 18–24.
- [57] Gross CG. Locomotor activity following lateral frontal lesions in rhesus monkeys. *J Comp Physiol Psychol* 1963, 56: 232–236.

- [58] Gross CG, Weiskrantz L. Some changes in behavior produced by lateral frontal lesions in the macaque. In: Warren JM, and Akert K (Eds.). *The Frontal Granular Cortex and Behavior*. New York: McGraw-Hill, 1964: 74–101.
- [59] Schachar RJ, Logan GD. Impulsivity and inhibitory control in normal development and childhood psychopathology. *Dev Psychol* 1990.
- [60] van der Meere J, Vreeling HJ, Sergeant J. A motor presetting study in hyperactive, learning disabled and control children. *J Child Psychol Psychiatry* 1992, 33: 1347–1354.
- [61] Durston S, Tottenham NT, Thomas KM, Davidson MC, Eigsti IM, Yang YH, *et al.* Differential patterns of striatal activation in young children with and without ADHD. *Biol Psychiatry* 2003, 53: 871–878.
- [62] Schulz KP, Fan J, Tang CY, Newcorn JH, Buchsbaum MS, Cheung AM, *et al.* Response inhibition in adolescents diagnosed with attention deficit hyperactivity disorder during childhood: an event-related fMRI study. *Am J Psychiatry* 2004, 161: 1650–1657.
- [63] Levy F. The dopamine theory of attention deficit hyperactivity disorder (ADHD). *Aust N Z J Psychiatry* 1991, 25: 277–283.
- [64] Arnsten AF, Li BM. Neurobiology of executive functions: catecholamine influences on prefrontal cortical functions. *Biol Psychiatry* 2005, 57: 1377–1384.
- [65] Halperin JM, Schulz KP. Revisiting the role of the prefrontal cortex in the pathophysiology of attention-deficit/hyperactivity disorder. *Psychol Bull* 2006, 132: 560–581.
- [66] Arnsten AF. Toward a new understanding of attention-deficit hyperactivity disorder pathophysiology: an important role for prefrontal cortex dysfunction. *CNS Drugs* 2009, 23 Suppl 1: 33–41.
- [67] Krause KH, Dresel SH, Krause J, la Fougere C, Ackenheil M. The dopamine transporter and neuroimaging in attention deficit hyperactivity disorder. *Neurosci Biobehav Rev* 2003, 27: 605–613.
- [68] Lynch CJ, Steer ML. Evidence for high and low affinity alpha 2-receptors. Comparison of [3H]norepinephrine and [3H] phentolamine binding to human platelet membranes. *J Biol Chem* 1981, 256: 3298–3303.
- [69] Hein L, Altman JD, Kobilka BK. Two functionally distinct alpha(2)-adrenergic receptors regulate sympathetic neurotransmission. *Nature* 1999, 402: 181–184.
- [70] Wang M, Ji JZ, Li BM. The alpha(2A)-adrenergic agonist guanfacine improves visuomotor associative learning in monkeys. *Neuropsychopharmacology* 2004, 29: 86–92.
- [71] Kopeckova M, Pacit I, Goetz P. Polymorphisms of dopamine-beta-hydroxylase in ADHD children. *Folia Biol (Praha)* 2006, 52: 194–201.
- [72] Comings DE, Gade-Andavolu R, Gonzalez N, Blake H, Wu S, MacMurray JP. Additive effect of three noradrenergic genes (ADRA2a, ADRA2C, DBH) on attention-deficit hyperactivity disorder and learning disabilities in Tourette syndrome subjects. *Clin Genet* 1999, 55: 160–172.
- [73] Arnsten AFT, Steere JC, Hunt RD. The contribution of α_2 - noradrenergic mechanisms to prefrontal cortical cognitive function: Potential significance to attention deficit hyperactivity disorder. *Arch Gen Psychiatry* 1996, 53: 448–455.
- [74] Steere JC, Arnsten AF. The α_2 -noradrenergic receptor agonist guanfacine improves visual object discrimination reversal performance in aged rhesus monkeys. *Behav Neurosci* 1997, 111: 883–891.
- [75] Wang M, Tang ZX, Li BM. Enhanced visuomotor associative learning following stimulation of alpha(2A)-adrenoceptors in the ventral prefrontal cortex in monkeys. *Brain Res* 2004, 1024: 176–182.
- [76] Wang M, Ramos BP, Paspalas CD, Shu Y, Simen A, Duque A, *et al.* Alpha2A-adrenoceptors strengthen working memory networks by inhibiting cAMP-HCN channel signaling in prefrontal cortex. *Cell* 2007, 129: 397–410.
- [77] Ramos BP, Stark D, Verduzco L, van Dyck CH, Arnsten AFT. alpha 2A-adrenoceptor stimulation improves prefrontal cortical regulation of behavior through inhibition of cAMP signaling in aging animals. *Learn Mem* 2006, 13: 770–776.
- [78] Ji XH, Ji JZ, Zhang H, Li BM. Stimulation of alpha2-adrenoceptors suppresses excitatory synaptic transmission in the medial prefrontal cortex of rat. *Neuropsychopharmacology* 2008, 33: 2263–2271.
- [79] Sagvolden T, Russell VA, Aase H, Johansen EB, Farshbaf M. Rodent models of attention-deficit/hyperactivity disorder. *Biol Psychiatry* 2005, 57: 1239–1247.
- [80] Russell VA. Overview of animal models of attention deficit hyperactivity disorder (ADHD). *Curr Protoc Neurosci* 2011, Chapter 9: Unit9. 35.
- [81] Okamoto K, Aoki K. Development of a strain of spontaneously hypertensive rats. *Jpn Circ J* 1963, 27: 282–293.
- [82] Sagvolden T. Behavioral validation of the spontaneously hypertensive rat (SHR) as an animal model of attention-deficit/hyperactivity disorder (AD/HD). *Neurosci Biobehav Rev* 2000, 24: 31–39.
- [83] Knardahl S, Sagvolden T. Open-field behavior of spontaneously hypertensive rats. *Behav Neural Biol* 1979, 27: 187–200.
- [84] Wultz B, Sagvolden T, Moser EI, Moser MB. The spontaneously hypertensive rat as an animal model of attention-deficit hyperactivity disorder: effects of methylphenidate on exploratory behavior. *Behav Neural Biol* 1990, 53: 88–102.
- [85] Sagvolden T, Pettersen MB, Larsen MC. Spontaneously hypertensive rats (SHR) as a putative animal model of

- childhood hyperkinesis: SHR behavior compared to four other rat strains. *Physiol Behav* 1993, 54: 1047–1055.
- [86] Sagvolden T, Slatta K, Arntzen E. Low doses of methylphenidate (Ritalin) may alter the delay-of-reinforcement gradient. *Psychopharmacology (Berl)* 1988, 95: 303–312.
- [87] Meneses A, Perez-Garcia G, Ponce-Lopez T, Tellez R, Gallegos-Cari A, Castillo C. Spontaneously hypertensive rat (SHR) as an animal model for ADHD: a short overview. *Rev Neurosci* 2011, 22: 365–371.
- [88] Berger DF, Sagvolden T. Sex differences in operant discrimination behaviour in an animal model of attention-deficit hyperactivity disorder. *Behav Brain Res* 1998, 94: 73–82.
- [89] Lou HC, Henriksen L, Bruhn P, Borner H, Nielsen JB. Striatal dysfunction in attention deficit and hyperkinetic disorder. *Arch Neurol* 1989, 46: 48–52.
- [90] Jesmin S, Sakuma I, Togashi H, Yamaguchi T, Ueno K, Yoshioka M, *et al.* Altered expression of endothelin and its receptors in the brain of SHR-SP at malignant hypertensive stage. *J Cardiovasc Pharmacol* 2004, 44 Suppl 1: S11–15.
- [91] Jesmin S, Togashi H, Mowa CN, Ueno K, Yamaguchi T, Shibayama A, *et al.* Characterization of regional cerebral blood flow and expression of angiogenic growth factors in the frontal cortex of juvenile male SHRSP and SHR. *Brain Res* 2004, 1030: 172–182.
- [92] Chappell PB, Riddle MA, Scahill L, Lynch KA, Schultz R, Arnsten A, *et al.* Guanfacine treatment of comorbid attention-deficit hyperactivity disorder and Tourette's syndrome: Preliminary clinical experience. *J Am Acad Child Adolesc Psychiatry* 1995, 34: 1140–1146.
- [93] Horrigan JP, Barnhill LJ. Guanfacine for treatment of attention-deficit hyperactivity disorder in boys. *J Child Adolesc Psychopharmacol* 1995, 5: 215–223.
- [94] Hunt RD, Arnsten AF, Asbell MD. An open trial of guanfacine in the treatment of attention-deficit hyperactivity disorder. *J Am Acad Child Adolesc Psychiatry* 1995, 34: 50–54.
- [95] Scahill L, Chappell PB, Kim YS, Schultz RT, Katsovich L, Shepherd E, *et al.* A placebo-controlled study of guanfacine in the treatment of children with tic disorders and attention deficit hyperactivity disorder. *Am J Psychiatry* 2001, 158: 1067–1074.
- [96] Findling RL, McBurnett K, White C, Youcha S. Guanfacine extended release adjunctive to a psychostimulant in the treatment of comorbid oppositional symptoms in children and adolescents with attention-deficit/hyperactivity disorder. *J Child Adolesc Psychopharmacol* 2014, 24: 245–252.
- [97] Connor DF, Arnsten AF, Pearson GS, Greco GF. Guanfacine extended release for the treatment of attention-deficit/hyperactivity disorder in children and adolescents. *Expert Opin Pharmacother* 2014, 15: 1601–1610.
- [98] Bymaster FP, Katner JS, Nelson DL, Hemrick-Luecke SK, Threlkeld PG, Heiligenstein JH, *et al.* Atomoxetine increases extracellular levels of norepinephrine and dopamine in prefrontal cortex of rat: A potential mechanism for efficacy in attention deficit/hyperactivity disorder. *Neuropsychopharmacology* 2002, 27: 699–711.
- [99] Wang Y, Zheng Y, Du Y, Song DH, Shin YJ, Cho SC, *et al.* Atomoxetine versus methylphenidate in paediatric outpatients with attention deficit hyperactivity disorder: a randomized, double-blind comparison trial. *Aust N Z J Psychiatry* 2007, 41: 222–230.
- [100] Muller U, Clark L, Lam ML, Moore RM, Murphy CL, Richmond NK, *et al.* Lack of effects of guanfacine on executive and memory functions in healthy male volunteers. *Psychopharmacology (Berl)* 2005, 182: 205–213.
- [101] Jerie P. Clinical experience with guanfacine in long-term treatment of hypertension. Part II: adverse reactions to guanfacine. *Br J Clin Pharmacol* 1980, 10 Suppl 1: 157S–164S.