

Reduced middle cingulate gyrus volume in late-onset schizophrenia in a Chinese Han population: a voxel-based structural MRI study

Dear Editor:

Numerous magnetic resonance imaging (MRI) studies have demonstrated that patients with early-onset schizophrenia (EOS) have widespread structural abnormalities in the cortical gray matter^[1], suggesting that neurobiological processes play a central role in the structural abnormalities underlying the pathophysiology of schizophrenia^[2]. In addition, volumetric abnormalities have been used to identify individuals at risk of mental states of psychosis by assessing patterns of individual whole-brain neuroanatomical abnormalities^[3]. However, few studies have investigated the abnormalities of brain structure in late-onset schizophrenia (LOS, age at onset >40 years), a distinct subtype characterized by prominent hallucinations and delusions, using voxel-based morphometry (VBM) analysis^[4]. However, to the best of our knowledge, there is no relevant report in the Chinese Han population. Thus, investigation of these abnormalities in LOS in this population could provide neuroanatomical clues for LOS. In the current study, we used VBM to analyze the gray matter volumes in Chinese Han patients with LOS and determined whether gray matter volume is associated with the psychotic symptoms, duration of illness, and medication dosage.

This study was approved by the Ethics Committee of the Second Xiangya Hospital of Central South University. The entire study design and procedures involved were in accordance with the Declaration of Helsinki. Risks and benefits of the study were presented in detail, and all participants gave written informed consent. Gray matter volume was measured in 20 patients with LOS and 17 well-matched healthy controls using VBM analysis (see supplementary material for Materials and Methods). Detailed demographics of patients with LOS and control subjects had been reported previously^[5].

The VBM analysis showed a significantly lower volume

of the middle cingulate gyrus in patients with LOS than in healthy controls (topological FDR corrected, $P < 0.05$, Fig. 1, Supplementary Table 1). In contrast, no region in the gray matter was significantly higher in LOS patients. Previous MRI studies in patients with LOS primarily focused on particular brain areas using classical measurements based on regions of interest (ROIs), in addition to the ventricles, periventricular white matter, and subcortical nuclei^[6]. VBM analysis is more sensitive than ROI analysis to structural changes of the whole brain. So, our VBM findings are more detailed and compatible with previous studies reporting cortical atrophy in patients with LOS^[6]. However, our VBM findings were not consistent with a previous report from Japan^[7]. This may be attributed to differences in the patients' age (range 40–58 years vs 41–73 years) and duration of illness (average duration 2.9 years vs 5.2 years)^[7], as well as the anatomical variations in different ethnic




Fig. 1. Regions with lower gray matter volume in late-onset schizophrenia than in healthy controls (topological FDR corrected $P < 0.05$). Late-onset schizophrenia patients, $n = 20$; healthy controls, $n = 17$.

populations^[8]. Also, a lower cingulate volume has been reported in elderly patients with schizophrenia^[9]. Abnormal levels of fractional anisotropy in the middle cingulum have been correlated with positive symptoms in antipsychotic-naïve first-episode patients with schizophrenia^[10]. Therefore, the middle cingulate gyrus may be the primary area affected in LOS and may play a crucial role in its pathology.

No correlation was found between the decrease in gray matter volume in the middle cingulate gyrus and the severity of psychiatric symptoms, illness duration, or antipsychotic dose (two-tailed, $P < 0.05$). Therefore, the current findings suggest that changes in gray matter volume are evident in the typical population of patients with LOS, and volumetric deficits in the middle cingulate gyrus may indicate susceptibility to schizophrenia.

ACKNOWLEDGEMENTS

The work was supported by the National Natural Science Foundation of China (30900486, 81371480, 81271484, 81471361, 81100996, and 81200838), and the National Basic Research Development Program (973) of China (2012CB517904). J.T. was supported by the Sheng-Hua Yuying Project of Central South University, China.

Hong Li¹, Jinsong Tang¹, Liping Chen², Yanhui Liao¹, Bing Zhou³, Ying He¹, Zongchang Li¹, Luxian Lv⁴, Yi Zeng⁵, Xiaogang Chen^{1,6,7}

¹*Institute of Mental Health, the Second Xiangya Hospital, Central South University, Changsha 410011, China*

²*The Dalian 7th People's Hospital, Dalian 116023, China*

³*Department of Radiology, the Second Xiangya Hospital, Central South University, Changsha 410011, China*

⁴*Department of Psychiatry, Henan Mental Hospital, The Second Affiliated Hospital of Xinxiang Medical University, Xinxiang 453002, China*

⁵*Department of Geriatrics, the Second Xiangya Hospital, Central South University, Changsha 410011, China*

⁶*Key Laboratory of Psychiatry and Mental Health of Hunan Province, Changsha, China*

⁷*National Technology Institute of Psychiatry, Changsha 410011, China*

Corresponding authors: Jinsong Tang, Xiaogang Chen and Liping Chen. E-mail: tangjinsonghn@gmail.com,

chenxghn@163.com, clp-lcc@163.com

© Shanghai Institutes for Biological Sciences, CAS and Springer-Verlag Berlin Heidelberg 2015

REFERENCES

- [1] Ellison-Wright I, Glahn DC, Laird AR, Thelen SM, Bullmore E. The anatomy of first-episode and chronic schizophrenia: an anatomical likelihood estimation meta-analysis. *Am J Psychiatry* 2008, 165: 1015–1023.
- [2] Brent BK, Thermenos HW, Keshavan MS, Seidman LJ. Gray matter alterations in schizophrenia high-risk youth and early-onset schizophrenia: a review of structural MRI findings. *Child Adolesc Psychiatr Clin N Am* 2013, 22: 689–714.
- [3] Koutsouleris N, Meisenzahl EM, Davatzikos C, Bottlender R, Frodl T, Scheuerecker J, *et al.* Use of neuroanatomical pattern classification to identify subjects in at-risk mental states of psychosis and predict disease transition. *Arch Gen Psychiatry* 2009, 66: 700–712.
- [4] Vahia IV, Palmer BW, Depp C, Fellows I, Golshan S, Kraemer HC, *et al.* Is late-onset schizophrenia a subtype of schizophrenia? *Acta Psychiatr Scand* 2010, 122: 414–426.
- [5] Chen L, Chen X, Liu W, Wang Q, Jiang T, Wang J, *et al.* White matter microstructural abnormalities in patients with late-onset schizophrenia identified by a voxel-based diffusion tensor imaging. *Psychiatry Res* 2013, 212: 201–207.
- [6] Lesser IM, Miller BL, Swartz JR, Boone KB, Mehlinger CM, Mena I. Brain imaging in late-life schizophrenia and related psychoses. *Schizophr Bull* 1993, 19: 773–782.
- [7] Egashira K, Matsuo K, Mihara T, Nakano M, Nakashima M, Watanuki T, *et al.* Different and shared brain volume abnormalities in late- and early-onset schizophrenia. *Neuropsychobiology* 2014, 70: 142–151.
- [8] Bai J, Abdul-Rahman MF, Rifkin-Graboi A, Chong YS, Kwek K, Saw SM, *et al.* Population differences in brain morphology and microstructure among Chinese, Malay, and Indian neonates. *PLoS One* 2012, 7: e47816.
- [9] Frisoni GB, Prestia A, Adorni A, Rasser PE, Cotelli M, Soricelli A, *et al.* *In vivo* neuropathology of cortical changes in elderly persons with schizophrenia. *Biol Psychiatry* 2009, 66: 578–585.
- [10] Cheung V, Chiu CP, Law CW, Cheung C, Hui CL, Chan KK, *et al.* Positive symptoms and white matter microstructure in never-medicated first episode schizophrenia. *Psychol Med* 2011, 41: 1709–1719.

Received date: 2015-01-08; Accepted date: 2015-04-02

(Supplementary material is available in the online version of this article at <http://dx.doi.org/10.1007/s12264-015-1525-1>.)