**RESEARCH HIGHLIGHT** 

## The Other Face of the Nucleus Accumbens: Aversion

Ying Han<sup>1,2</sup> · Lin Lu<sup>1,2,3</sup>

Received: 18 April 2016/Accepted: 30 May 2016/Published online: 8 August 2016 © Shanghai Institutes for Biological Sciences, CAS and Springer Science+Business Media Singapore 2016

The nucleus accumbens (NAc) is thought to integrate information that is conveyed by (1) dopaminergic inputs from the midbrain and (2) glutamatergic inputs from limbic and cortical regions, including the amygdala, prefrontal cortex (PFC), hippocampus, and thalamus [1], and signal to the basal ganglia motor system to guide appropriate behaviors. Addictive drug use is proposed to hijack this system, enhancing the brain's reactivity to drug cues and triggering drug-seeking and relapse after prolonged withdrawal [2, 3]. Both positive reinforcement (rewarding effects of drugs) and negative reinforcement (aversive emotional state associated with withdrawal) have been considered to play critical roles in the etiology and maintenance of drug addiction, and NAc neurons are capable of processing both reward and aversion [4, 5]. Optogenetic activation of inputs from the PFC, ventral hippocampus (vHipp), and basolateral amygdala (BLA) to the NAc drives positive reinforcement and facilitates rewardseeking behavior [6, 7], but a recent paper in *Nature* [8] has identified another face of the NAc. The pathway from the paraventricular nucleus of the thalamus (PVT) to the NAc is key to the aversive emotional state of drug withdrawal.

Lin Lu linlu@bjmu.edu.cn

Glutamatergic synaptic transmission within the NAc has been recognized as a primary target for addictive drugs to produce adaptive synaptic changes and modulate behavioral output. Cocaine selectively increases the presynaptic release probability of excitatory synapses within the PFC-NAc pathway but not the BLA-NAc pathway, and amygdala fibers have a relatively low probability of transmitter release [7, 9]. Chronic non-contingent or contingent cocaine exposure and withdrawal evoke input- and cell type-specific plasticity in the NAc, which may underlie behavioral adaptations in addiction, such as craving and relapse [7, 10]. Optogenetic-mediated, pathway-specific stimulation reveals that the activation of inputs from the PFC, vHipp, and BLA to the NAc is rewarding and can reinforce instrumental behavior [6, 7]. However, still unknown is the specific NAc circuitry that underlies negative emotional and motivational states after withdrawal.

Using retrograde neuronal tracing, Zhu et al. [8] showed that the NAc is densely innervated by the PVT, in addition to its well-characterized inputs from the PFC, vHipp, and BLA. Brief light stimulation of fibers that project from the PVT to the NAc evoked robust α-amino-3-hydroxy-5-methyl-4-isoxazolepropionic acid receptor (AMPAR)-mediated excitatory postsynaptic currents in medium spiny neurons (MSNs), indicating that the PVT is a source of glutamatergic afferents to the NAc. Behaviorally, optogenetically activating the PVT-NAc pathway evokes avoidance of a light-paired chamber in the realtime place preference assay, which depends on glutamatergic but not dopaminergic transmission in the NAc. Thus, in contrast to other inputs to the NAc, the PVT input mediates aversion rather than reward. This suggests the possibility that this pathway is a specific neuronal circuit involved in the negative emotional state associated with drug withdrawal.



<sup>&</sup>lt;sup>1</sup> Institute of Mental Health, National Clinical Research Center for Mental Disorders, Key Laboratory of Mental Health, and Peking University Sixth Hospital, Peking University, Beijing 100191, China

<sup>&</sup>lt;sup>2</sup> National Institute on Drug Dependence and Beijing Key Laboratory of Drug Dependence, Peking University, Beijing 100191, China

<sup>&</sup>lt;sup>3</sup> Peking-Tsinghua Center for Life Sciences and PKU-IDG/ McGovern Institute for Brain Research, Peking University, Beijing 100871, China



Fig. 1 Synaptic changes in the paraventricular nucleus of the thalamus-to-nucleus accumbens pathway after opiate withdrawal. *Left* glutamatergic fibers that originate in the paraventricular nucleus of the thalamus (PVT) innervate medium spiny neurons (MSNs) in the nucleus accumbens (NAc) and control motivated behaviors. *Middle* chronic morphine treatment selectively strengthens

Zhu et al. [8] used optogenetic and chemogenetic approaches to selectively silence the PVT-NAc pathway and test the effects of this manipulation on opiate withdrawal-induced somatic signs and place aversion. Robust Fos expression in PVT neurons that project to the NAc was precipitated by naloxone and induced by spontaneous withdrawal. Optogenetic inhibition of PVT axons expressing archaerhodopsin-3 in the NAc during naloxoneprecipitated withdrawal suppressed physical signs and avoidance of the withdrawal chamber in morphinedependent mice but not in drug-naive mice. Using inhibitory designer receptors exclusively activated by designer drugs, silencing the PVT-NAc pathway by intra-NAc injection of clozapine-N-oxide before four conditioned place aversion (CPA) training sessions prevented the formation of aversive memory.

However, foot-shock and LiCl injection also increased Fos expression in PVT neurons that project to the NAc. Silencing this pathway during conditioning reduced the expression of foot-shock- and LiCl-induced CPA, indicating that the PVT-NAc pathway also mediates aversion induced by these stimuli. Together, it is speculated that the PVT-to-NAc circuit plays important roles in the modulation and encoding of general aversion such as stress and fear conditioning.

MSNs can be classified into two groups based on the dopamine receptors that they express ( $D_1$  or  $D_2$  [D1R or D2R]). These subtypes have different projection targets

 $PVT \rightarrow D_2$  receptor (D2R)-MSN synapses *via* the insertion of GluA2-lacking Ca<sup>2+</sup>-permeable AMPARs (CP-AMPARs), triggering strong physical signs and robust aversive memory. *Right* depotentiation of the PVT input onto D2R-MSNs by optogenetic induction of long-term depression (LTD) restores normal transmission at these synapses and suppresses aversive withdrawal symptoms.

(direct or indirect to the midbrain) and exert complementary and sometimes opposing actions on behaviors that are controlled by the corticostriatal system [11-13]. Previous studies have reported an increase in synaptic strength in D1R-MSNs but not D2R-MSNs after cocaine withdrawal, most likely through a postsynaptic mechanism [10, 14]. Zhu et al. [8] showed that chronic morphine treatment increased the AMPAR/N-methyl-D-aspartate receptor ratio and the insertion of GluA2-lacking Ca<sup>2+</sup>-permeable AMPARs at  $PVT \rightarrow D2R$ -MSN synapses but not  $PVT \rightarrow D1R$ -MSN synapses. Moreover, they used an established in vivo long-term depression (LTD) protocol to reduce synaptic transmission at  $PVT \rightarrow D2R$ -MSN synapses (i.e., light pulses at 1 Hz for 15 min). This optogenetic protocol restored normal transmission at  $PVT \rightarrow D2R$ -MSN synapses, without influencing synaptic strength at PVT  $\rightarrow$  D1R-MSN synapses, and ultimately reduced the expression of opiate withdrawal symptoms and CPA of the withdrawal chamber.

Previous studies have shown that orexin/hypocretin transmission in the PVT primarily originating from the lateral hypothalamus plays an important role in drug-seeking behavior [15, 16]. Besides the NAc, the PVT also specifically projects to the ventral tegmental area, PFC, and hippocampus, which are widely implicated in reward-related behaviors [16]. Further exploration of the specific neuronal subpopulations and circuits in the PVT mediating reward-seeking and aversive symptoms seems warranted.

The study by Zhu *et al.* [8] is a landmark in our understanding of the neuronal circuitry that underlies opiate withdrawal and highlights the role of plasticity at  $PVT \rightarrow D2R$ -MSN synapses in the negative emotional and motivational states associated with opiate withdrawal (Fig. 1). Their results suggest that some of the neuronal adaptations associated with opiate dependence may be reversible. Optogenetic protocols that target specific synapses and use specific stimulation parameters to cause potentiation or de-potentiation may inspire novel treatments for opiate addiction. In humans, novel deep brain stimulation or transcranial magnetic stimulation protocols may restore normal synaptic transmission, thus relieving withdrawal symptoms [17].

## References

- Sesack SR, Grace AA. Cortico-basal ganglia reward network: microcircuitry. Neuropsychopharmacology 2010, 35: 27–47.
- Dong Y, Nestler EJ. The neural rejuvenation hypothesis of cocaine addiction. Trends Pharmacol Sci 2014, 35: 374–383.
- Volkow ND, Morales M. The brain on drugs: from reward to addiction. Cell 2015, 162: 712–725.
- 4. Wise RA, Koob GF. The development and maintenance of drug addiction. Neuropsychopharmacology 2014, 39: 254–262.
- Wenzel JM, Rauscher NA, Cheer JF, Oleson EB. A role for phasic dopamine release within the nucleus accumbens in encoding aversion: a review of the neurochemical literature. ACS Chem Neurosci 2015, 6: 16–26.
- Stuber GD, Sparta DR, Stamatakis AM, van Leeuwen WA, Hardjoprajitno JE, Cho S, et al. Excitatory transmission from the amygdala to nucleus accumbens facilitates reward seeking. Nature 2011, 475: 377–380.

- Britt JP, Benaliouad F, McDevitt RA, Stuber GD, Wise RA, Bonci A. Synaptic and behavioral profile of multiple glutamatergic inputs to the nucleus accumbens. Neuron 2012, 76: 790–803.
- Zhu Y, Wienecke CF, Nachtrab G, Chen X. A thalamic input to the nucleus accumbens mediates opiate dependence. Nature 2016, 530: 219–222.
- Suska A, Lee BR, Huang YH, Dong Y, Schluter OM. Selective presynaptic enhancement of the prefrontal cortex to nucleus accumbens pathway by cocaine. Proc Natl Acad Sci USA 2013, 110: 713–718.
- Pascoli V, Terrier J, Espallergues J, Valjent E, O'Connor EC, Luscher C. Contrasting forms of cocaine-evoked plasticity control components of relapse. Nature 2014, 509: 459–464.
- Gerfen CR, Surmeier DJ. Modulation of striatal projection systems by dopamine. Annu Rev Neurosci 2011, 34: 441–466.
- Ferguson SM, Eskenazi D, Ishikawa M, Wanat MJ, Phillips PE, Dong Y, et al. Transient neuronal inhibition reveals opposing roles of indirect and direct pathways in sensitization. Nat Neurosci 2011, 14: 22–24.
- Kravitz AV, Tye LD, Kreitzer AC. Distinct roles for direct and indirect pathway striatal neurons in reinforcement. Nat Neurosci 2012, 15: 816–818.
- Pascoli V, Turiault M, Luscher C. Reversal of cocaine-evoked synaptic potentiation resets drug-induced adaptive behaviour. Nature 2012, 481: 71–75.
- 15. Matzeu A, Zamora-Martinez ER, Martin-Fardon R. The paraventricular nucleus of the thalamus is recruited by both natural rewards and drugs of abuse: recent evidence of a pivotal role for orexin/hypocretin signaling in this thalamic nucleus in drugseeking behavior. Front Behav Neurosci 2014, 8: 117.
- Martin-Fardon R, Boutrel B. Orexin/hypocretin (Orx/Hcrt) transmission and drug-seeking behavior: is the paraventricular nucleus of the thalamus (PVT) part of the drug seeking circuitry? Front Behav Neurosci 2012, 6: 75.
- Luigjes J, van den Brink W, Feenstra M, van den Munckhof P, Schuurman PR, Schippers R, et al. Deep brain stimulation in addiction: a review of potential brain targets. Mol Psychiatry 2012, 17: 572–583.